




PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C12N 15/86, 9/12, C07K 14/18, A61K 48/00, A01K 67/027, C12N 5/10, 7/04		A1	(11) International Publication Number: WO 99/50432
(21) International Application Number: PCT/IB99/00523		(43) International Publication Date: 7 October 1999 (07.10.99)	
(22) International Filing Date: 25 March 1999 (25.03.99)			
(30) Priority Data: 60/079,562 27 March 1998 (27.03.98) US			
(71) Applicant (for all designated States except US): CYTOS BIOTECHNOLOGY AG [CH/CH]; Einsteinstrasse, CH-8093 Zurich (CH).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(71)(72) Applicants and Inventors: RENNER, Wolfgang, A. [CH/CH]; Weinbergstrasse 64, CH-8006 Zurich (CH). NIEBA, Lars [DE/CH]; Gottfried-Keller Strasse 63B, CH-8400 Winterthur (CH). BOORSMA, Marco [NL/NL]; Telemannstraat 10d, NL-8915 CB Leeu Warden (NL).		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	

(54) Title: INDUCIBLE ALPHAVIRAL GENE EXPRESSION SYSTEM



(57) Abstract

The present invention provides novel expression vectors which permit tight regulation of gene expression in eucaryotic cells. More specifically, the invention provides DNA vectors comprising nucleotide sequences that are transcribed to form RNA molecules which are then replicated by a temperature-sensitive replicase to form additional RNA molecules. The RNA molecules produced by replication contain a nucleotide sequence which may be translated to produce a protein of interest or which encode one or more untranslated RNA molecules. Also provided are methods for producing heterologous proteins and untranslated RNA molecules. Further provided are methods for administering heterologous proteins and untranslated RNA molecules to individuals. In addition, pharmaceutical compositions are provided comprising the DNA and RNA molecules of the invention and a pharmaceutically acceptable carrier.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INDUCIBLE ALPHAVIRAL GENE EXPRESSION SYSTEM

Background of the Invention

Field of the Invention

5 The present invention relates to novel expression vectors which permit tight regulation of gene expression in eucaryotic cells. The invention also relates to methods for producing proteins and RNA molecules and methods for administering proteins and RNA molecules to a plant or animal.

Related Art

10 The ability to precisely control the expression of genes introduced into animal or human cells, or in whole organisms, will enable significant progress in many areas of biology and medicine. For instance, methods that allow the intentional manipulation of gene expression will facilitate the analysis of genes whose expression cannot be tolerated constitutively or at a certain stage of development. These methods will also be valuable for clinical applications such as gene therapy, where the expression of a therapeutic gene must be regulated in
15 accordance with the needs of the patient.

To be of broad benefit, gene regulation techniques must allow for rapid, robust, precise and reversible induction of gene activity. As reviewed in Saez, E. *et al.*, (*Curr. Opin. Biotechnol.* 8:608-616 (1997)), an ideal system should fulfill
20 the following requirements:

1. Specificity -- The system must be indifferent to endogenous factors and activated only by exogenous stimuli.
2. Non-interference -- The components of the system should not affect unintended cellular pathways.
- 25 3. Inducibility -- In the inactive state, the basal activity of the system should be minimal, while in the active state high levels of gene expression should be rapidly inducible.

-2-

4. Bioavailability of the inducer -- Inducing stimuli should rapidly penetrate to the site of interest.
5. Reversibility -- Inducing stimuli should clear swiftly to allow the system to rapidly return to the inactive state.

5 Early methods for controlling gene expression in mammals were based on endogenous elements, such as cytokine response elements or heat-shock proteins. Due to a high level of basal expression in the uninduced state, and pleiotropic effects brought about by general inducing agents, these systems lacked the specificity required to regulate genes in mammalian cells and organisms.

10 More advance schemes have sought to avoid these problems by constructing switching mechanisms that rely on non-mammalian elements. The fundamental principle of these systems is based on the existence of a small molecule (the inducer) that modifies the activity of a synthetic transcription factor which regulates the expression of the target gene through a heterologous promoter. Increased specificity is achieved by selecting inducers that do not
15 affect mammalian physiology, and by assembling chimeric transactivators with minimal homology to natural transcription factors which do not interact with endogenous mammalian promoters.

20 The most common system currently in use for the regulation of gene expression is the tetracycline-based system (Gossen and Bujard, *Proc. Natl. Acad. Sci. USA* 89:5547 (1992)). This system is based on the continuous expression of a fusion protein where the tetracycline repressor protein (tetR) is converted into an activator by fusion to the transcriptional activation domain of the VP16 protein. In the absence of tetracycline, this chimeric tetracycline transactivator (tTA) activates gene expression through binding to a multimer of the natural tetR
25 binding site (tetO) placed upstream of a minimal promoter. In the presence of tetracycline, the tTA undergoes a conformational change that prevents it from binding to the tetO site, thereby arresting expression of the target gene. Because of its significant advantages over the existing approaches, the tTA system is highly useful for inducible gene expression and this system has been successfully
30 used for the production of a number of proteins (Wimmel *et al.*, *Oncogene* 9:995 (1994); Fröh *et al.*, *EMBO J.* 13:3236 (1994); Yu *et al.*, *J. Virol.* 70:4530 (1996)).

However, serious problems resulting from the toxicity of the tTA protein have been reported with the tTA system, and several cell types have been shown to be unable to tolerate expression of the tTA protein (Schocket *et al.*, *Proc. Natl. Acad. Sci. USA* 92:6522 (1995); Howe *et al.*, *J. Biol. Chem.* 23:14168 (1995); Schocket and Schatz, *Proc. Natl. Acad. Sci. USA* 93:5173 (1996); Bohl *et al.*, *Nat. Med.* 3:299 (1997)). While the toxicity of tTA in cultured cells encumbers the establishment of stable clones with proper tetracycline regulation, this tTA toxicity is a more significant problem in gene therapy and may prevent the use of the tTA system in gene therapy altogether.

A further problem of the tTA system is its notable degree of basal expression. Basal expression can result from the activation of the reporter constructs in the absence of bound transactivator, and/or the inability of tetracycline to completely quell tTA transactivation. High basal expression limits the inducibility of the system, and prevents the conductance of experiments with highly toxic proteins (Furth *et al.*, *Proc. Natl. Acad. Sci. USA* 91:9302 (1994); Hennighausen *et al.*, *J. Cell. Biochem.* 59:463 (1995) Kistner *et al.*, *Proc. Natl. Acad. Sci. USA* 93:10933 (1996); Hoffmann *et al.*, *Nucleic Acids Res.* 25:1078 (1997)).

In the case of stable clones or transgenic animals, some of this basal expression can be attributed to interference from chromosomal regions into which the foreign DNA integrates. While all inducible systems are equally susceptible to integration effects, it is possible that the basal activity of the tTA system is due to the fact that this system requires the constant presence of tetracycline to efficiently suppress transcription, something that may not always be attainable, particularly *in vivo*. Basal expression and the requirement that tetracycline be present to suppress gene expression are reasons why the tTA system is not used in gene therapy.

Two gene control systems based on components of mammalian steroid hormone receptors are known (Saez, E. *et al.*, *Curr. Opin. Biotechnol.* 8:608-616 (1997)). Both combine a truncated form of the progesterone receptor hormone-binding domain with the yeast GAL4 DNA-binding moiety, and the transactivation domain of VP16 protein. The mutated progesterone receptor

-4-

moiety fails to bind progesterone, but it retains the ability to bind the progesterone and glucocorticoid antagonist mifepristone (RU486), such that, in the presence of RU486, the fusion protein (called GVLP or TAXI) activates transcription through a multimer of the GAL4 DNA binding site placed upstream of a minimal promoter.

An important advantage of the systems described immediately above is that they appear to have more favorable kinetics than tetracycline approaches because lipophilic hormones are quickly metabolized and have short half-lives *in vivo*. Further, such hormones may also penetrate less accessible tissues more efficiently than tetracycline. However, the main disadvantage of the hormone receptor systems is their very high level of basal expression. In transient and stable transfections of various cell types, a high level of basal activity dampens the inducibility of these approaches, resulting in induction ratios that are rarely over 20-fold (Wang *et al.*, *Proc. Natl. Acad. Sci. USA* 91:8180 (1994); Mangelsdorf *et al.*, *Cell* 83:835 (1995); Wang *et al.*, *Nat. Biotech.* 15:239 (1997)).

Another approach to regulating gene expression relies on a method of inducing protein dimerization derived from studies on the mechanism of action of immunosuppressive agents (Saez, E. *et al.*, *Curr. Opin. Biotechnol.* 8:608-616 (1997)). Using a synthetic homodimer of FK506, a general strategy was devised to bring together any two peptides simply by endowing them with the domain of FKBP12 to which FK506 binds. Since immunosuppressive drugs, such as cyclosporin A or rapamycin must be used in this approach, the *in vivo* application of this protein dimerization approach is very limited.

All of the above mentioned strategies regulate expression by controlling the level of transcription of mRNA. Since this mRNA transcription mechanism is always influenced to some extent by the chromosomal region into which the foreign DNA is inserted, precise regulation fails due to the lack of control over the integration mechanism. Although techniques are available for the site-specific insertion of DNA by homologous recombination, insertion frequencies are far too low to allow this strategy to succeed for gene therapy on a general basis.

-5-

Another gene expression system is based on alphaviruses (Lundstrom, K., *Curr. Opin. Biotechnol.* 8:578-582 (1997)). Several members of the alphavirus family, Sindbis (Xiong, C. *et al.*, *Science* 243:1188-1191 (1989); Schlesinger, S., *Trends Biotechnol.* 11:18-22 (1993)), SFV (Liljeström, P. & Garoff, H., *Bio/Technology* 9:1356-1361 (1991)) and others (Davis, N.L. *et al.*, *Virology* 171:189-204 (1989)), have received considerable attention for the use as virus-based expression vectors for a variety of different proteins (Lundstrom, K., *Curr. Opin. Biotechnol.* 8:578-582 (1997); Liljeström, P., *Curr. Opin. Biotechnol.* 5:495-500 (1994)).

Alphaviruses are positive stranded RNA viruses which replicate their genomic RNA entirely in the cytoplasm of the infected cell and without a DNA intermediate (Strauss, J. and Strauss, E., *Microbiol. Rev.* 58:491-562 (1994)). The concept that alphaviruses can be developed as expression vectors was first established nearly ten years ago (Xiong, C. *et al.*, *Science* 243:1188-1191 (1989)). Since then, several improvements have made the use of these RNA replicons as expression vectors more practical (Lundstrom, K., *Curr. Opin. Biotechnol.* 8:578-582 (1997)).

DNA vectors have been developed for both Sindbis (Herweijer, H. *et al.*, *Hum. Gene Ther.* 6:1495-1501 (1995); Dubensky, T.W. *et al.*, *J. Virol.* 70:508-519 (1996)) and SFV (Berglund, P. *et al.*, *Trends Biotechnol.* 14:130-134 (1996)). Eukaryotic promoters are introduced in these vectors upstream from the alphavirus replicase gene (consisting of the four non-structural protein genes (nsP1-4)) which are translated as one or two polyproteins which are then proteolytically cleaved (Strauss, J. and Strauss, E., *Microbiol. Rev.* 58:491-562 (1994)). DNA is transcribed to RNA from the recombinant eukaryotic promoter in the nucleus and transported to the cytoplasm, where the replicase catalyzes the replication of the alphavirus RNA molecule as during normal replication of the alphavirus RNA molecule (Strauss, J. and Strauss, E., *Microbiol. Rev.* 58:491-562 (1994)). Only transient expression of heterologous sequences has been possible until recently due to the cytopathogenicity of the alphavirus replicase (Lundstrom, K., *Curr. Opin. Biotechnol.* 8:578-582 (1997)).

-6-

About 20 years ago Weiss *et al.* (Weiss, B. *et al.*, *J. Virol.* 33:463-474 (1980)) established a persistently infected culture of BHK cells. The mutation responsible for this phenotype has been recently identified (Dryga, S.A. *et al.*, *Virology* 228:74-83 (1997)). Another mutation allowing the regulation of the mRNA transcription via temperature shifts was identified by Burge and Pfefferkorn (Burge, B.W. & Pfefferkorn, E.R., *Virology* 30:203-214 (1966)) and described in more detail by Xiong *et al.* (Xiong, C. *et al.*, *Science* 243:1188-1191 (1989)).

Vectors containing alphaviral sequences have been developed which show promise for use in DNA immunizations (Hariharan, M. *et al.*, *J. Virol.* 72:950-958 (1998)), ribozyme expression (Smith S. *et al.*, *J. Virol.* 71:9713-9721 (1997)), and *in vivo* expression of heterologous proteins in mammalian tissues (Altman-Hamamdzie S. *et al.*, *Gene Ther.* 4:815-822 (1997)).

Summary of the Invention

The present invention provides compositions and methods for regulated expression of proteins or untranslated RNA molecules in recombinant host cells. More specifically, the present invention provides polynucleotides and methods which allow precise regulation of the amount of specific RNA molecules produced in stably transfected recombinant host cells. This precise regulation results from the use of a temperature-sensitive RNA-dependent RNA polymerase (*i.e.*, a replicase) which only replicates RNA molecules, to form new RNA molecules, at permissive temperatures.

In one general aspect, the DNA expression vectors of the invention comprise a 5' promoter which is capable of initiating transcription *in vivo*, 5' and/or 3' sequences enabling replication of the RNA molecule (*cis*-acting sequence elements), and a subgenomic promoter 5' to the gene of interest, as well as a sequence of interest which is translatable only after one or more RNA-dependent RNA replication events. These RNA-dependent RNA replication events are catalyzed by a regulatable RNA-dependent RNA

-7-

polymerase which may be encoded by the same mRNA molecule that is produced by transcription of the DNA vector or by a different mRNA molecule.

In another aspect, the invention provides DNA molecules comprising polynucleotides which encode RNA molecules comprising (a) at least one *cis*-acting sequence element, (b) a first open reading frame having a nucleotide sequence encoding a non-cytopathic, temperature-sensitive RNA-dependent RNA polymerase, and (c) at least one second nucleotide sequence which encodes one of the following:

(i) a second open reading frame encoding a protein, or portion thereof, wherein the second open reading frame is in a translatable format after one or more RNA-dependent RNA replication events;

(ii) a sequence complementary to all or part of the second open reading frame of (i); and

(iii) a sequence encoding an untranslated RNA molecule (e.g., an antisense RNA molecule, tRNA molecule, rRNA molecule, or ribozyme), or complement thereof.

The invention further provides single- and multiple-vector systems for expressing a second nucleotide sequence described above. In a single-vector system, sequences encoding the first open reading frame and the second nucleotide sequence are present on the same nucleic acid molecule. In a multiple-vector system, sequences encoding the first open reading frame, or sub-portions thereof, and the second nucleotide sequence are present on one or more separate nucleic acid molecules.

When sequences encoding the first and second open reading frame are present either on the same nucleic acid molecule or in the same vector (*i.e.*, in a single-vector system), a region will be present 5' to the second open reading frame which inhibits translation of this open reading frame.

The temperature-sensitive replicase may be "cold" or "hot" sensitive and thus will only efficiently catalyze RNA-dependent RNA replication at temperatures either above or below the restrictive temperature. In one embodiment, the DNA molecules of the invention encode an RNA-dependent

-8-

RNA polymerase that has replicase activity at temperatures below 34°C and has low or undetectable replicase activity at temperatures of 34°C and above.

Further provided are RNA transcription products of the DNA molecules of the invention and alphaviral particles containing packaged RNA molecules of the invention. When packaged RNA molecules are produced, the second open reading frame may encode one or more proteins required for such packaging (e.g., Sindbis structural proteins).

In another aspect, the nucleic acid molecules of the invention encode one or more cytokine, lymphokine, tumor necrosis factor, interferon, toxic protein, prodrug converting enzyme, or other protein.

In yet another aspect, the nucleic acid molecules of the invention encode an untranslated RNA molecule, such as an antisense RNA molecule, tRNA molecule, rRNA molecule, or ribozyme.

The invention also provides methods for making recombinant host cells comprising introducing nucleic acid molecules of the invention into host cells. Further provided are recombinant host cells produced by the introduction of nucleic acid molecules of the invention. In one embodiment, some or all of these recombinant host cells contain one or more DNA molecules of the invention which are stably maintained.

The invention further provides the pCYTs vector of SEQ ID NO:1, as well as isolated nucleic acid molecules comprising polynucleotides having the nucleotide sequence of SEQ ID NO:1.

The present invention also provides methods for producing proteins and untranslated RNA molecules in recombinant host cells comprising growing host cells under suitable culture conditions, introducing nucleic acid molecules of the invention into host cells, and recovering the proteins or untranslated RNA molecules produced by the recombinant host cells.

Methods are also provided for the regulated expression of heterologous polypeptides, including cytokines, lymphokines, tumor necrosis factors, interferons, toxic proteins, and prodrug converting enzymes.

Further provided are proteins and untranslated RNA molecules produced by the methods of the invention.

-9-

The invention also provides methods for regulating the expression of heterologous proteins in recombinant host cells comprising growing host cells under suitable culture conditions, introducing nucleic acid molecules of the invention into the host cells, and changing the temperature of the host cell culture from either a permissive temperature to a restrictive temperature or a restrictive temperature to a permissive temperature. In one embodiment, the nucleic acid molecules of the invention are introduced into prokaryotic or eukaryotic host cells which are then cultured *in vitro*. In related embodiments, these host cells are cultured in a serum-free or protein-free medium.

Additionally provided are methods for producing proteins in recombinant host cells comprising growing host cells under suitable culture conditions, infecting said host cells with alphaviral particles containing RNA molecules of the invention, and recovering the protein.

Also provided are methods for the introduction and expression of nucleic acid molecules of the invention in recombinant host cells within an individual. When these recombinant host cells are intended to express polypeptide or untranslated RNA sequences in an individual, the nucleic acid molecules of the present invention may be introduced into host cells either *in vivo* or *ex vivo*. When the nucleic acid molecules are introduced into host cells *ex vivo*, the recombinant host cells can either be administered to the individual from which they were obtained or to a different individual. In certain embodiments, the host cells are mammalian keratinocytes, epithelial cells, or fibroblasts which are reintroduced into the same mammal from which they were obtained.

The invention further provides methods for regulating the expression of proteins or untranslated RNA molecules in individuals comprising administering nucleic acid molecules of the invention to individuals and changing the temperature of at least a portion of these individuals from either a permissive temperature to a restrictive temperature or a restrictive temperature to a permissive temperature.

The invention also provides methods for administering proteins and untranslated RNA molecules to individuals comprising administering nucleic acid molecules of the invention to individuals and changing the temperature of at least

-10-

a portion of the individuals from a restrictive temperature to a permissive temperature.

The invention additionally provides methods for regulating the expression of proteins and untranslated RNA molecules in individuals comprising administering recombinant host cells of the invention to these individuals and changing the temperature of at least a portion of these individuals from either a permissive temperature to a restrictive temperature or a restrictive temperature to a permissive temperature.

In one embodiment, the host cells are obtained from the same individual into which the recombinant host cells are administered. In another embodiment, the recombinant host cells are keratinocytes.

The present invention also provides pharmaceutical compositions comprising nucleic acid molecules of the invention and a pharmaceutically acceptable carrier.

The present invention further provides genetically engineered, non-human animals which contain nucleic acid molecules of the invention in at least some of their cells. Also provided are genetically engineered, non-human animals which contain DNA molecules of the invention stably integrated into the genome of some or all the animal's cells. The invention also provides methods for producing genetically engineered, non-human animals comprising introducing cells containing nucleic acid molecules of the invention into these animals, introducing nucleic acid molecules of the invention into the cells of these animals *in vivo*, or introducing DNA molecules of the invention into germ line cells to produce transgenic animals containing the sequence of interest in their somatic and germ line cells.

Brief Description of the Figures

FIG. 1. The DNA of pCYTs (1) is inserted into the nucleus. The eukaryotic promoter (solid horizontal arrow) drives transcription (2) into mRNA (3). Translation (4) of the first open reading frame (ORF) of the mRNA results in the production of a temperature-sensitive replicase (*ts*-replicase protein) (5).

-11-

The second open ORF encoding the gene of interest is not accessible to ribosomes. Thus no translation (6) of the gene of interest occurs. At low temperature the *ts*-replicase catalyzes replication (7) of the mRNA (3) into full-length (-) strand RNA (8). The *ts*-replicase also catalyzes subsequent replications (9, 10) into full-length (+) strand RNA (11) and subgenomic RNA (12). The subgenomic RNA (12) is then translated (13) into the protein of interest (not shown). The combination of amplification and qualitative change of the RNA results in unprecedented tightness and regulatability of the expression of the gene of interest.

Abbreviations in FIG. 1 are as follows: Rous Sarcoma Virus promoter (RSV pr.), *cis*-acting sequence elements (CSE), non-structural proteins 1-4 (nsP1, nsP2, nsP3, nsP4), gene of interest (G.O.I.), and subgenomic promoter (S.G.)

FIG. 2 is a schematic representation of the pCYTts vector. The pCYTts vector contains, in addition to the elements shown in FIG. 1, an ampicillin resistance marker for selection in bacterial cells and a ColE1 sequence which directs high copy number bacterial amplification. The pCYTts vector was prepared as described in Example 1.

FIG. 3A-3D shows the complete cDNA sequence of pCYTts (SEQ ID NO:1).

FIG. 4A-4B. GFP (FIG. 4A) and SEAP (FIG. 4B) production at different temperatures. Cells stably transfected with pCYTtsGFP or pCYTtsSEAP were grown for 48 hours at the indicated temperatures (closed diamonds and open squares). Two independent experiments are shown in each of FIG. 4A and FIG. 4B. GFP fluorescence (FIG. 4A) was determined as described in Example 2. SEAP activity (FIG. 4B) was determined colorimetrically as described in Example 3. The maximal protein expression was determined for both proteins to be 29°C. The activities of the proteins were calculated relative to the maximal value at 29°C.

FIG. 5A-5B. Time course of GFP (FIG. 5A) and SEAP (FIG. 5B) production in BHK cells stably transfected with pCYTtsGFP (FIG. 5A) and pCYTtsSEAP (FIG. 5B) at 30°C. GFP production was measured by spectrofluorophotometry and quantified in fluorescence units per 10⁶ cells, as

-12-

described in Example 2. SEAP production per 10^6 cells was determined as described in Example 3 by measuring enzymatic activity using p-nitrophenylphosphate as a substrate. The amount of SEAP produced after 80 hours was estimated to be over 10^7 molecules per cell.

5 FIG. 6A-6B. The start of GFP (FIG. 6A) or SEAP (FIG. 6B) mRNA transcription in a mixed population of BHK cells stably transfected with pCYTsGFP or pCYTsSEAP was determined by measuring the amount of GFP or SEAP produced (see Examples 2 and 3). Cells were incubated for 2, 4, 6, 8 and 10 hours at 29°C (black boxes), and then grown for another 24 hours at 37°C (open boxes).

10 FIG. 7A-7B. These figures show the results of the experiments obtained using stable pCYTsGFP transfected-BHK cells which were transiently transfected with a plasmid coding for the structural proteins of Sindbis virus. The conditions of the experiments were as follows: (I) incubation phase of transfected BHK cells at 29°C or 37°C for 48 hours; (II) the supernatant of the cells was put onto a new BHK cell layer and the cells were shifted to the indicated temperatures; (III) incubation at either 29°C or 37°C of the new BHK cell layer for 6 hours; and (IV) washing of the cells and final incubation at 29°C or 37°C for 48 hours. Infection events were visualized by the expression of the marker gene GFP as described in Example 2. FIG. 7A shows the result of two separate experiments performed as described above. Fluorescence indicates GFP expression, whereas no fluorescence indicates no detectable GFP expression. FIG. 7B shows the results of four separate experiments performed as described above. The + and - symbols indicate whether GFP expression was detected.

20 FIG. 8A-8B. Western blot of β -interferon (β -INF) (FIG. 8A) and erythropoietin (EPO) (FIG. 8B) expressed in the pCYTs system. FIG. 8A shows in lane 1 marker, lane 2 positive control, lane 3 supernatant of β -INF expressing cells following incubation at 37°C for 3 days, lane 4 supernatant of β -INF expressing BHK cells (transient transfection) following incubation at 29°C for 3 days, lane 5 supernatant of GFP expressing BHK cells, lane 6 marker, lane 7 supernatant of β -INF expressing BHK cells (mixed population) following incubation at 29°C for 3 days, and lane 8 marker. FIG. 8B shows in lane 1

-13-

marker. lane 2 supernatant of EPO expressing cells (transient transfection) following incubation at 29°C for 5 days, lane 3 supernatant of EPO expressing cells following incubation at 37°C for 5 days, lane 4 supernatant of GFP expressing BHK cells. and lane 5 marker.

5 FIG. 9 shows a Western Blot of EPO. The samples in each lane are as follows: lane 1 EPO standard; lane 2 supernatant of stably pCYTs504-Epo transfected cells at 37°C for 4 days; lane 3 supernatant of stably pCYTs504-Epo transfected cells at 29°C for 4 days; lane 4 marker.

10 FIG. 10 shows a dot blot of EPO. Spot + shows EPO standard. spots 2 and 10 supernatant of BHK cells (2), spot 3 GFP expressing BHK cells incubated at 30°C, spot 4 1C4 cells incubated at 37°C, spot 8 supernatant of the BHK cells infected with CYTs504Epo RNA containing viral particles incubated at 37°C after 2 days, spot 9 supernatant of BHK cell infected with CYTs504Epo RNA containing viral particles incubated at 30°C for 2 days.

15 FIG. 11 shows an overview of one embodiment of the invention. This embodiment is directed to the production of recombinant human EPO using host cell infected with packaged RNA replicons produced by baby hamster kidney (BHK) cell line 1C4/4. This BHK cell line was produced as described below in Example 6.

20 FIG. 12 shows a flow chart of one embodiment of the invention. According to the process described in this figure. RNA replicons are produced by the stable EPO-packaging BHK cell line 1C4/4 and isolated from the culture medium. Wild-type BHK cells, which may be cultured in either serum- or protein-free culture media, are then infected with the replicons. The protein produced by the infected cells is then purified by conventional processes. The process outlined in this figure can be readily scaled up for production of large quantities of human EPO or other proteins.

25

Detailed Description of the Preferred Embodiments

The present invention is directed to improved expression vectors that are regulatable and non-cytopathic, as well as methods for using these vectors to produce proteins and RNA molecules of interest.

5 The invention provides polynucleotides and methods which allow the precise regulation of the amount of specific RNA molecules produced in host cells. This precise regulation results from the use of a temperature-sensitive RNA-dependent RNA polymerase which will only replicate RNA molecules, to form additional RNA molecules, at permissive temperatures.

10 The invention is further directed to inducible gene expression systems employing alphavirus DNA vectors to create stable cell lines carrying genes encoding a non-cytopathic, temperature-sensitive, viral non-structural replicase protein. For example, the activity of the temperature-sensitive replicase used in the Examples, set out below, is switched on by reducing the temperature of the transfected cells from a temperature of 37°C to a temperature lower than 34°C. Host cell expression at 37°C is below the level of detection and the induction profile is independent of the chromosomal integration site.

Definitions

20 The following definitions are provided to clarify the subject matter which the inventors consider to be the present invention.

As used herein, the term "alphavirus" refers to any of the RNA viruses included within the genus *Alphavirus*. Descriptions of the members of this genus are contained in Strauss and Strauss, *Microbiol. Rev.*, 58:491-562 (1994). Examples of alphaviruses include Aura virus, Bebaru virus, Cabassou virus, Chikungunya virus, Easter equine encephalomyelitis virus, Fort morgan virus, Getah virus, Kyzylagach virus, Mayoaro virus, Middleburg virus, Mucambo virus, Ndumu virus, Pixuna virus, Tonate virus, Trinita virus, Una virus, Western equine encephalomyelitis virus, Whataroa virus, Sindbis virus (SIN), Semliki forest virus (SFV), Venezuelan equine encephalomyelitis virus (VEE), and Ross River virus.

-15-

As used herein, when the term "purified" is used in reference to a molecule, it means that the concentration of the molecule being purified has been increased relative to molecules associated with it in its natural environment. Naturally associated molecules include proteins, nucleic acids, lipids and sugars but generally do not include water, buffers, and reagents added to maintain the integrity or facilitate the purification of the molecule being purified. For example, even if mRNA is diluted with an aqueous solvent during oligo dT column chromatography, mRNA molecules are purified by this chromatography if naturally associated nucleic acids and other biological molecules do not bind to the column and are separated from the subject mRNA molecules.

As used herein, when the term "isolated" is used in reference to a molecule, the term means that the molecule has been removed from its native environment. For example, a polynucleotide or a polypeptide naturally present in a living animal is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated." Further, recombinant DNA molecules contained in a vector are considered isolated for the purposes of the present invention. Isolated RNA molecules include *in vivo* or *in vitro* RNA replication products of DNA and RNA molecules. Isolated nucleic acid molecules further include synthetically produced molecules. Additionally, vector molecules contained in recombinant host cells are also isolated. Thus, not all "isolated" molecules need be "purified."

As used herein, the phrase "low or undetectable," when used in reference to gene expression level, refers to a level of expression which is either significantly lower than that seen when the gene is maximally induced (*e.g.*, at least five fold lower) or is not readily detectable by the methods used in the following examples section.

As used herein, the phrase "individual" refers to multicellular organisms and includes both plants and animals. Preferred multicellular organisms are animals, more preferred are vertebrates, even more preferred are mammals, and most preferred are humans.

As used herein, the phrase "*cis*-acting" sequence refers to nucleic acid sequences to which a replicase binds to catalyze the RNA-dependent replication

-16-

of RNA molecules. These replication events result in the replication of the full-length and partial RNA molecules and, thus, the alphavirus subgenomic promoter is also a "cis-acting" sequence. *Cis*-acting sequences may be located at or near the 5' end, 3' end, or both ends of a nucleic acid molecule, as well as internally.

5 As used herein, the phrase "RNA-Dependent RNA polymerase" refers to a polymerase which catalyzes the production of an RNA molecule from another RNA molecule. This term is used herein synonymously with the term "replicase."

 As used herein, the phrase "non-infective packaged RNA molecules" refers to packaged RNA molecules which will essentially undergo only one round of host cell infection and are not pathogenic. These molecules are thus
10 "infective" but only for a single infectious entry into a host cell and are not capable of reproducing to form additional infectious particles.

 As used herein, the term "transcription" refers to the production of RNA molecules from DNA templates catalyzed by RNA polymerases.

15 As used herein, the phrase "RNA-dependent RNA replication event" refers to processes which result in the formation of an RNA molecule using an RNA molecule as a template.

 As used herein, the term "vector" refers to an agent (*e.g.*, a plasmid or virus) used to transmit genetic material to a host cell. A vector may be composed
20 of either DNA or RNA.

 As used herein, the term "heterologous sequence" refers to a second nucleotide sequence present in a vector of the invention. The term "heterologous sequence" also refers to any amino acid or RNA sequence encoded by a heterologous DNA sequence contained in a vector of the invention. Heterologous
25 nucleotide sequences can encode proteins or RNA molecules normally expressed in the cell type in which they are present or molecules not normally expressed therein (*e.g.*, Sindbis structural proteins).

 As used herein, the phrase "untranslated RNA" refers to an RNA sequence or molecule which does not encode an open reading frame or encodes an open
30 reading frame, or portion thereof, but in a format in which an amino acid sequence will not be produced (*e.g.*, no initiation codon is present). Examples of such molecules are tRNA molecules, rRNA molecules, and ribozymes. Antisense

-17-

RNA may be untranslated but, in some instances (see Example 11), antisense sequences can be converted to a translatable sense strand from which a polypeptide is produced.

As used herein the phrase "gene therapy" refers to the transfer of heterologous genetic information into cells for the therapeutic treatment of diseases or disorders. The heterologous nucleotide sequence is transferred into a cell and is expressed to produce a polypeptide or untranslated RNA molecule.

As used herein, the phrase "temperature-sensitive" refers to an enzyme which readily catalyzes a reaction at one temperature but catalyzes the same reaction slowly or not at all at another temperature. An example of a temperature-sensitive enzyme is the replicase protein encoded by the pCYTts vector, which has readily detectable replicase activity at temperatures below 34°C and has low or undetectable activity at 37°C.

As used herein, the phrase "permissive temperature" refers to temperatures at which an enzyme has relatively high levels of catalytic activity.

As used herein, the phrase "restrictive temperature" refers to temperatures at which an enzyme has low or undetectable levels of catalytic activity. Both "hot" and "cold" sensitive mutants are known and, thus, a restrictive temperature may be higher or lower than a permissive temperature.

As used herein, the term "recombinant host cell" refers to a host cell into which one or more nucleic acid molecules of the invention have been introduced.

When the terms "one," "a," or "an" are used in this disclosure, they mean "at least one" or "one or more," unless otherwise indicated.

Alphaviral Vectors of the Invention

The DNA vectors of the present invention are constitutively transcribed in host cells to produce mRNA molecules having two open reading frames. These open reading frames, which may or may not be produced from the same nucleic acid molecule, encode a temperature-sensitive replicase and a heterologous gene of interest. The first open reading frame is translated to produce a temperature-dependent RNA-dependent RNA polymerase. The second

-18-

open reading frame, encoding all or part of one or more polypeptides of interest, is not translated until after at least one RNA-dependent RNA replication event.

The DNA expression vectors comprise a 5' promoter which is capable of initiating synthesis of RNA *in vivo*. 5' and/or 3' sequences enabling replication of the RNA molecule (5' and 3' *cis* acting sequence elements), as well as a sequence of interest which is translatable only after at least one replication event. Replication is catalyzed by a regulatable RNA-dependent RNA polymerase which is encoded alternatively on the same or on a different mRNA molecule. The sequence of interest may be encoded in sense, plus (+) orientation downstream of a viral RNA promoter. Translation of the coding sequence of the gene of interest is inhibited by a 5' sequence which, in the case of the single-vector system, will generally be the replicase sequence. In the multiple-vector system, a 5' sequence can inhibit translation by having one or more short open reading frames with associated stop codons which lead to the detachment of ribosomes. Similarly, any sequence which inhibits the traveling or binding of ribosomes to the sequence of interest can be used as a 5' sequence which inhibits translation (Voet and Voet, BIOCHEMISTRY, John Wiley & Sons, Inc. (1990)).

Another method for preventing translation of nucleotide sequences in most biological systems involves the insertion of the sequence in an antisense direction. This method of inhibiting translation is based on the principle that translation will generally only occur after the replication of this minus (-) strand RNA into a plus strand having an open reading frame in a sense orientation. The translated sense strand is formed by RNA replication and serves as a template for ribosomes and protein synthesis. As shown in Example 11, production of amino acid sequences can occur even when the gene of interest is inserted into the DNA molecule in an orientation which will result in the formation of antisense RNA sequence 3' to the subgenomic promoter. Thus, the second open reading frame may also comprise a sequence complementary to all or part of the second open reading frame, described above and expression of the encoded amino acid sequence will still occur. When the production of an untranslated antisense RNA sequence is desired, the RNA molecule can be designed so that it will not serve as a template for protein synthesis. For example, the RNA can be designed so

-19-

that an initiation codon is not present.

Untranslated antisense RNA molecules can be used to inhibit translation of mRNA expressed in recombinant host cells. The use of antisense nucleic acid molecules to regulate gene expression is known in the art (*see, e.g., Kawamata, H. et al., Br. J. Cancer* 77:71-78 (1998); Bechler, K., *Biochem. Biophys. Res. Commun.* 241:193-199 (1997); Urakami, S. et al., *Biochem. Biophys. Res. Commun.* 241:24-30 (1997)) and the use of the present vectors to deliver such molecules to host cells is within the scope of the invention.

In addition, instead of a second open reading frame, RNA molecules directly produced by transcription of a DNA sequence of the invention may encode RNA sequences which are neither translated nor present in an antisense orientation. Examples of such untranslated RNA molecules include tRNA molecules, rRNA molecules, and ribozymes. A considerable number of ribozyme sequences with defined catalytic activities are known in the art (*see, e.g., Brown, J., Nucleic Acids Res.* 26:353-354 (1998); Xie, Y. et al., *Proc. Natl. Acad. Sci. USA* 94:13777-13781 (1997); Lavrovsky, Y. et al., *Biochem. Mol. Med.* 62:11-22 (1997); Chapman, K. and Szostak, J., *Chem. Biol.* 2:325-333 (1995)). Further, ribozymes have been used to "knockout" the expression of a specific gene in eucaryotic cells as part of a ribozyme-mediated, message deletion strategy (Xie, Y. et al., *Proc. Natl. Acad. Sci. USA* 94:13777-13781 (1997)). Additionally, alphaviral replicons have been used to express a functional ribozyme in mammalian cells (Smith S. et al., *J. Virol.* 71:9713-9721 (1997)). The regulated expression of such ribozymes, and other untranslated RNA molecules, is thus within the scope of the present invention.

The invention is exemplified by the schematic diagram shown in FIG. 1. These embodiments of the invention are directed to DNA vectors which are transcribed to produce a mRNA molecule having two open reading frames, which encode a replicase and a gene of interest. The DNA vectors contain a promoter sequence which drives transcription of these vectors to produce mRNA molecules having coding sequences of both open reading frames. The mRNA sequences of the first open reading frame are translated to produce a replicase required for the expression of the RNA sequences of the second open reading frame. The second

-20-

open reading frame encodes one or more proteins of interest.

Further, once the first mRNA molecule has been transcribed from the DNA vector, additional RNA-dependent RNA replication events can occur to amplify the first mRNA sequence and to produce RNA molecules with strand polarity which is the opposite of the first mRNA sequence.

As shown in FIG. 1, sections (7)-(8), (10), and (12)-(13), the second open reading frame of a DNA molecules of the invention will only be expressed after partial replication of a full-length RNA molecule. This partial replication of the full-length RNA molecules is driven by a promoter sequence composed of RNA (*e.g.*, an alphaviral subgenomic promoter sequence).

While the gene of interest may be encoded by the same RNA molecule as the replicase protein, this gene may also be encoded by a separate RNA molecule. Thus, the invention further provides both single- and multiple-vectors systems for expressing a gene of interest.

In a single-vector system of the invention, sequences encoding the first open reading frame and the second nucleotide sequence are components of the same nucleic acid molecule. Thus, all of the components required for regulated expression of the gene of interest are contained in a single nucleic acid molecule (*i.e.*, DNA or RNA).

In a multiple-vector system of the invention, sequences encoding the first open reading frame, or sub-portions thereof, and the second nucleotide sequence are components of different nucleic acid molecules. These multiple-vector systems thus may comprise two or more nucleic acid molecules. For example, nsP2, nsP4, and the gene of interest can each be encoded by different DNA vectors. Further, one or more of these DNA vectors can be designed to stably integrate into the host cell genome. When expression of a gene of interest is desired in a cell type containing one or more stably integrated DNA molecules of the invention, expression of the gene of interest will require the introduction of nucleic acid molecules (DNA or RNA) encoding the components of the system into the cells not present in the integrated molecule(s).

While any functional promoter can be used to drive the transcription of mRNA from the DNA vector, the promoter is preferably a constitutive RNA

-21-

polymerase II promoter (e.g., Rous Sarcoma Virus (RSV), cytomegalovirus (CMV), simian virus 40 (SV40), myeloproliferative sarcoma virus (MPSV), glucocorticoid, metallothionein, *Herpes simplex* virus thymidine kinase (HSVTK), human immuno deficiency (HIV), mouse mammary tumor virus (MMTV), human polyomavirus BK (BKV), or Moloney murine leukemia virus (MuLV) promoter). Additional promoters suitable for use in the practice of the present invention are known in the art (see, e.g., Lee, A. *et al.*, *Mol. Cells.* 7:495-501 (1997); Artuc, M. *et al.*, *Exp. Dermatol.* 4:317-321 (1995)).

The vector will generally also contain selection markers for cloning and amplification of the vector sequences in procaryotic and eucaryotic organisms. The pCYTts vector, for example, contains an ampicillin resistance marker for positive selection in bacterial host cells and an *E. coli* origin of replication (i.e., ColE1). A considerable number of sequences encoding additional selection markers and origins of replication are known in the art (see, e.g., Sambrook, J. *et al.*, eds., *MOLECULAR CLONING. A LABORATORY MANUAL*, 2nd. edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989); Ausubel, F. *et al.*, eds., *CURRENT PROTOCOLS IN MOLECULAR BIOLOGY*, John H. Wiley & Sons, Inc. (1997)).

The replicase protein coding sequences, the 5' and 3' *cis*-acting sequences (when present), and the junction sequences containing the subgenomic promoter will normally be derived from a virus, preferably from an alphavirus, most preferably from Sindbis virus.

When using alphavirus replicase proteins, in most instances, it is desirable to convert the cytopathic phenotype of the replicase protein to a non-cytopathic phenotype. Preferred mutations which confer such a phenotype are in the *nsp2* gene (e.g., the proline residue at position 726 is replaced with a serine residue). Mutations are known in the art which render the replicase protein non-cytopathic (Weiss *et al.*, *J. Virol.* 33:463-474 (1980); Dryga *et al.*, *Virology* 228:74-83 (1997)). These mutations may be introduced by a number of means, including site directed mutagenesis.

As noted above, when a non-cytopathic Sindbis virus replicase is used in the practice of the invention, a mutation may be introduced in the *nsp2* gene. One

-22-

such mutation results from the exchange of the proline residue at position 726 to another of the 20 natural occurring amino acids, such as a serine (abbreviated as "Pro 726 Ser"). Alternatively, any other mutation rendering the replicase molecule non-cytopathic is within the scope of the invention. The creation and the identification of mutations which render the Sindbis replicase non-cytopathic are described in more detail elsewhere (Weiss *et al.*, *J. Virol.* 33:463-474 (1980); Dryga *et al.*, *Virology* 228:74-83 (1997); patent application WO 97/38087). Further, methods for inducing such mutations are known in the art (*see, e.g.*, Sambrook, J. *et al.*, eds., *MOLECULAR CLONING, A LABORATORY MANUAL*, 2nd edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989); Ausubel, F. *et al.*, eds., *CURRENT PROTOCOLS IN MOLECULAR BIOLOGY*, John H. Wiley & Sons, Inc. (1997)).

Temperature sensitivity (*ts*) may be conferred, for example, by the introduction of a mutation in the *nsp4* gene of the replicase. Preferably, mutations which confer a temperature-sensitive phenotype upon replicase activities are in a protein in complementation group F (Lemm *et al.*, *J. Virol.* 64:3001-3011 (1990)). For example, a temperature-sensitive phenotype may be conferred by changing Gly 153 of *nsp4* to Glu. Additionally, any other mutation which renders replicase activity temperature-sensitive can be used in the practice of the invention. Methods for creating and identifying new temperature-sensitive mutants are described by Pfefferkorn (Burge and Pfefferkorn, *Virol.* 30:204-213 (1966); Burge and Pfefferkorn, *Virol.* 30:214-223 (1966)). Further, any method useful for producing and identifying *ts* mutants which allow for the temperature-sensitive regulation of replicase activity can be employed to generate and isolate such mutants.

While most temperature-sensitive mutants are "hot" sensitive, "cold" sensitive ones are also known (*see, e.g.*, Schwer, B. *et al.*, *Nucleic Acids Res.* 26:803-809 (1998), Mathe, E. *et al.*, *J. Cell Sci.* 111:887-896 (1998), Doedens, J. *et al.*, *J. Virol.* 71:9054-9064 (1997), Patterson, B. *et al.*, *J. Biol. Chem.* 272:27612-27617 (1997)). The temperature-sensitive replicase may be "cold" or "hot" sensitive and thus will catalyze RNA replication only at temperatures either above or below restrictive temperatures. In one embodiment, RNA replication

-23-

occurs at detectable levels only at temperatures lower than 34°C. In a related embodiment, the pCYTts vector, or variant thereof, is used to express an inserted gene of interest with expression being induced by reducing the temperature of cells containing the vector from 37°C to a temperature lower than about 34°C.

5 As shown in FIG. 4A-4B, permissive temperatures for the replicase encoded by the pCYTts vector are below about 34°C. Further, expression of the gene of interest increases when the temperature is increased from about 24°C until a maximal expression level is reached at about 29°C. Additionally, expression of the gene of interest increases as the temperature decreases from about 34°C.

10 Thus, permissive temperatures for the replicase activity encoded by the pCYTts vector are below 34°C, and include temperatures below 24°C, as well as 24°C, 25°C, 26°C, 27°C, 28°C, 29°C, 30°C, 31°C, 32°C, and 33°C and intervening fractional temperatures up to about 34°C.

In contrast to all previously known regulatable DNA expression systems,

15 the basal level of expression in recombinant host cells containing the pCYTts vector in the inactive state at 37°C is below the level of detection using standard methods (*e.g.*, those used in the following examples). This low level of expression is apparent from the data presented in FIG. 4A-4B, FIG. 8A-8B, FIG. 9, and FIG. 10. Further, the temperature-dependent induction profile of gene

20 expression appears to be independent of the chromosomal integration site and copy number.

In another embodiment, the sequence of interest and non-cytopathic, regulatable replicase (*e.g.*, nsp2 carrying the Pro 726 Ser mutation and nsp4 carrying the Gly 153 Glu mutation) are encoded by two separate DNA vectors.

25 In such an instance, the DNA vector carrying the sequence of interest carries both *cis*-acting sequences and a 5' region which inhibits translation of the sequence of interest. The non-cytopathic, regulatable replicase gene can also be encoded by a DNA molecule which is different than the one carrying the sequence of interest. Replication and translation of the sequence of interest in this multi-vectors system

30 is regulatable by temperature as in the one vector system.

The vectors of the invention can be also used to regulate the expression of more than one gene of interest. For example, recombinant host cells can be

-24-

transfected with more than one nucleic acid molecule of the invention wherein one nucleic acid molecule encodes both the replicase and a polypeptide of interest and additional nucleic acid molecules could encode additional polypeptides of interest. Similarly, when mutations conferring non-cytopathicity and temperature sensitivity are both used, genes encoding polypeptides having suitable mutations (e.g., Pro 726 Ser in nsp2 and Gly 153 Glu in nsp4) may be on separate nucleic acid molecules. Additional variations would be apparent to those skilled in the art.

As shown in Example 11, the sequence of interest can also be inserted in an antisense direction downstream from a functional promoter (e.g., the myeloproliferative sarcoma virus (MPSV) promoter). This construct is converted to a plus (+) strand with sense polarity as shown by the production of the protein of interest. These data demonstrate that antisense DNA fragments can be used with the present invention to express functional polypeptides, or subportions thereof. These data further indicate that the 5' and 3' CSEs may not be necessary for viral transcription when antisense DNA is used as a template for transcription.

The DNA molecules of the invention can also contain packaging signals which direct the packaging of RNA molecules into viral particles. These RNA molecules can be packaged in the presence of wild-type virus or defective helper virus RNA. A significant improvement was made with the development of defective helper RNA molecules (Bredenbeek, P. *et al.*, *J. Virol.* 67:6439-6446 (1993)). These RNA molecules contain *cis*-acting sequences, required for replication of the full-length transcription product, and subgenomic RNA promoter sequences which drive the expression of the structural protein genes. For example, in cells containing both RNA molecules with packaging signals and the defective helper virus RNA, alphaviral non-structural proteins allow for replication and amplification of the defective helper virus RNA sequences which are translated to produce virion structural proteins. Since the helper virus RNA lacks packaging signals, these molecules are not packaged into assembled virions. Thus, virion particles produced in this way contain essentially only RNA sequences encoding the gene of interest and, generally, other sequences required for temperature-sensitive regulation of gene expression. These non-infective

-25-

packaged RNA molecules do not contain sequences encoding virion structural proteins and, thus, undergo only one round of host cell infection and are not pathogenic.

Non-infective packaged RNA molecules can be used to infect a culture of suitable host cells simply by addition of the particles to culture medium containing these cells. The preparation of non-infective alphaviral particles is described in a number of sources, including "Sindbis Expression System", Version C, (*Invitrogen* Catalog No. K750-1).

One application of this system is directed to the temperature-dependent production of non-infective, packaged RNA molecules. These packaged RNA molecules may be produced by a number of means including using recombinant host cells containing two different DNA molecules (*e.g.*, a DNA molecule of the invention and a DNA molecule encoding a helper virus RNA sequence). For example, one of these DNA molecules will encode an RNA molecule which contains packaging signal sequences, sequences encoding a non-cytopathic, temperature-sensitive replicase, and the gene of interest. The other DNA molecule will contain sequences encoding alphaviral structural proteins downstream from an alphavirus subgenomic promoter. Using such a system, viral particles containing only RNA molecules with packaging signals will be produced at permissive temperatures in recombinant host cells. This is so because alphaviral structural proteins will only be produced at a permissive temperature. Additional variations of the above would be apparent to one skilled in the art.

A wide variety of nucleotide sequences of interest can be expressed by the gene expression system of the invention. These sequences include, but are not limited to, sequences encoding lymphokines, cytokines, toxins, enzymes, prodrug converting enzymes, antigens which stimulate immune responses, single chain antibodies, proteins which stimulate or inhibit immune responses, tumor necrosis factors, and various proteins with therapeutic applications (*e.g.*, growth hormones and regulatory factors).

As demonstrated in Example 5, heterologous sequences expressed by the vectors of the invention can also encode erythropoietin (EPO). EPO is a

-26-

glycoprotein which belongs to the cytokine family and induces terminal erythrocyte development. This protein also regulates red blood cell production.

Heterologous sequences can also encode a cytokine or lymphokine (e.g., β -interferon). Hematopoiesis is regulated by lymphokines and cytokines which stimulate the proliferation and/or differentiation of various hemopoietic cells. Representative examples of cytokines and lymphokines include interleukin-1 (IL-1), interleukin-2 (IL-2), interleukin-3 (IL-3), interleukin-4 (IL-4), interleukin-5 (IL-5), interleukin-6 (IL-6), interleukin-7 (IL-7), interleukin-8 (IL-8), interleukin-9 (IL-9), interleukin-10 (IL-10), interleukin-11 (IL-11), interleukin-12 (IL-12), interleukin-13 (IL-13), interleukin-14 (IL-14), interleukin-15 (IL-15), interleukin-16 (IL-16), interleukin-17 (IL-17), granulocyte colony stimulating factor (G-CSF), granulocyte-macrophage colony stimulating factor (GM-CSF), macrophage colony stimulating factor (M-CSF), and interferons.

Heterologous sequences can also encode secreted enzymes (e.g., secreted alkaline phosphatase), cytoplasmic enzymes (e.g., green fluorescent protein), or any number of other proteins with therapeutic applications (e.g., human insulin, human coagulation Factor VIII).

The vectors of the invention can also be used to express heterologous sequence encoding cytotoxic polypeptides. Cytotoxic polypeptides act to directly or indirectly inhibit cellular growth or metabolism. Representative examples of toxins include *Shigella* toxin, ricin, *Diphtheria* toxin, *Cholera* toxin, *Pseudomonas* exotoxin A, and *Herpes simplex* virus thymidine kinase (HSVTK). Within other embodiments of this invention, the heterologous sequence encodes a prodrug converting enzyme. A prodrug converting enzyme activates a compound with little or no cytotoxicity into a toxic product. Representative example are HSVTK, alkaline phosphatase, guanine phosphoribosyl transferase, and penicillin-V amidase. Examples of both cytotoxic polypeptides and prodrug converting enzymes are discussed in numerous sources including PCT/US97/06010, EP 0716148, and WO 96/17072.

Nucleotide sequences which may be used with the vectors of the invention include untranslated RNA molecules, such as antisense sequences, RNase P targeted sequences which induce gene down-regulation, and ribozymes. Smith

-27-

S. *et al.* (*J. Virol.* 71:9713-9721 (1997)) describes alphaviral vectors used to express ribozyme sequences.

The nucleic acid molecules of the invention can also be used to express virtually any protein, including ones which have not as yet been identified but are encoded by nucleotide sequences contained in, for example, cDNA libraries or host cell chromosomes. Example of such proteins include secreted proteins and proteins from various cellular compartments. Heterologous sequences expressed by the vectors of the invention can encode proteins and RNA molecules from non-human species (*e.g.*, other mammals, plants, fungi, bacteria or viruses). These heterologous sequence may further encodes viral membrane proteins (*e.g.*, HIV gp160) or viral polypeptides (*e.g.*, Sindbis structural proteins).

Sequences of the above described proteins may be readily obtained from a variety of sources, including for example the American Type Culture Collection (ATCC, Rockville, MD). Alternatively, cDNA sequences which encode the above-mentioned heterologous sequences may be obtained from cells which express such sequences. Methods for isolating both genomic and cDNA sequences encoding genes of interest are well known in the art (*see, e.g.*, Celis, J., ed., *CELL BIOLOGY*, Academic Press, 2nd edition, (1998); Sambrook, J. *et al.*, eds., *MOLECULAR CLONING, A LABORATORY MANUAL*, 2nd edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989); Ausubel, F. *et al.*, eds., *CURRENT PROTOCOLS IN MOLECULAR BIOLOGY*, John H. Wiley & Sons, Inc. (1997)). For example, mRNA can be isolated from a cell which expresses a sequence of interest, after which the sequence of interest is reverse transcribed with reverse transcriptase using oligo dT primers, random primers, specific primers, or combinations of each. The cDNA sequences may then be amplified by PCR using heat stable proof-reading polymerases. Alternatively, synthetic DNA sequences may be constructed and expressed with the vectors of the invention.

Nucleotide sequences may be added to the vectors of the invention which result in the production of a fusion protein. For example, such sequences can encode amino acids sequences which are fused to a protein encoded by a gene of interest and confer one or more functional characteristics upon the expression

-28-

product. These amino acid sequences include sequences which will target the gene product for export from the cell (e.g., a secretory sequence) or to a subcellular compartment (e.g., the nucleus). Such amino acid sequences further include sequences which facilitate purification (e.g., a six His "tag"). Depending on the amino acid sequence and the function imparted by the fused sequence, the added amino acid sequences may or may not be cleaved from the translation product.

Fusion proteins also include proteins which have domains or regions derived from various different proteins. Examples of such a fusion protein are those containing domain II of *Pseudomonas* exotoxin, a domain or amino acid sequence which has binding affinity for a cell surface receptor associated with a particular cell type, and another amino acid sequence with a preselected biological activity. Domain II of *Pseudomonas* exotoxin will translocate across cell membranes. Using this system, fusion proteins can be designed which will bind to specific cells types, will translocate across the cytoplasmic membranes of these cells, and will catalyze predetermined intracellular biological reactions. A system of this type is described in Pastan *et al.*, U.S. Patent No. 5,705,163. Methods for identifying amino acid sequences which bind to specific cell types are described in Wu, A., *Nature Biotech.* 14:429-431 (1996).

The vectors of the invention can also contain genetic elements which confer additional functional characteristics such as selection markers, sequences which result in high copy number host cell amplification, and sequences which allow for chromosomal integration of vector sequences.

Markers for the selection of prokaryotic and eukaryotic cells containing vectors the present invention are well known in the art and include tetracycline, ampicillin, neomycin, and kanamycin resistance. DNA molecules containing such sequences are available from numerous sources including Stratagene (11011 North Torrey Pines Road, La Jolla, CA 92037, USA) and Promega (2800 Woods Hollow Road, Madison, WI 53711, USA). Nucleotide sequences which result in high copy number amplification are also known in the art and include the ColEI sequence contained in the pCYTts vector.

Recombinant Host Cells

A variety of different recombinant host cells can be produced which contain the vectors of the invention. Alphaviruses are known to have a wide host range. Sindbis virus, for example, infects cultured mammalian, reptilian, and amphibian cells, as well as some insect cells (Clark, H., *J. Natl. Cancer Inst.* 51:645 (1973); Leake, C., *J. Gen. Virol.* 35:335 (1977); Stollar, V. in *THE TOGAVIRUSES*, R.W. Schlesinger, Ed., Academic Press. (1980), pp.583-621). Thus, numerous recombinant host cells can be used in the practice of the invention. BHK, COS, Vero, HeLa and CHO cells are particularly suitable for the production of heterologous proteins because they have the potential to glycosylate heterologous proteins in a manner similar to human cells (Watson, E. *et al.*, *Glycobiology* 4:227, (1994)) and can be selected (Zang, M. *et al.*, *Bio/Technology* 13:389 (1995)) or genetically engineered (Renner W. *et al.*, *Biotech. Bioeng.* 47:476 (1995); Lee K. *et al. Biotech. Bioeng.* 50:336 (1996)) to grow in serum-free medium, as well as in suspension.

When recombinant host cells capable of expressing a gene of interest are intended to be inserted into an individual, these cells will generally be from either another individual of the same genus and species or the same individual into which the cells will be inserted. Cells may be obtained from an individual by any number of means including surgical means and tissue biopsy.

Introduction of the polynucleotide vectors into host cells can be effected by methods described in standard laboratory manuals (*see, e.g.*, Sambrook, J. *et al.*, eds., *MOLECULAR CLONING, A LABORATORY MANUAL*, 2nd. edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989), Chapter 9; Ausubel, F. *et al.*, eds., *CURRENT PROTOCOLS IN MOLECULAR BIOLOGY*, John H. Wiley & Sons, Inc. (1997), Chapter 16). including methods such as electroporation, DEAE-dextran mediated transfection, transfection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction, and infection. Methods for the introduction of exogenous DNA sequences into host cells are discussed in Felgner, P. *et al.*, U.S. Patent No. 5,580,859.

Non-infective packaged RNA sequences can also be used to infect host

-30-

cells. These packaged RNA sequences can be introduced to host cells by adding them to the culture medium.

As noted *supra*, the vectors of the invention may also contain genetic elements which allow for chromosomal integration of vector sequences. Such elements are useful for the stable maintenance of heterologous sequences and include sequences which confer both site-specific and site-independent integration. Site-specific integration (*e.g.*, homologous integration) and site-independent integration, sometimes referred to as "random integration" can be used to introduce heterologous sequences of interest into eucaryotic chromosomes. Descriptions and methods for inserting genetic material into eucaryotic chromosomes are available from numerous sources including Sambrook, J. *et al.*, eds. (MOLECULAR CLONING, A LABORATORY MANUAL, 2nd. edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989)).

Production of Polypeptides and RNA Molecules

The vectors and recombinant host cells of the invention may be used for the production of polypeptides and RNA molecules. Thus, the invention provides methods for the regulated expression of polypeptides or RNA molecules in host cells, comprising the step of introducing nucleic acid sequences of the present invention into host cells and regulating the temperature to either repress or induce the production of RNA molecules encoding sequences of interest.

Recombinant host cells which express a gene of interest will generally either express this gene in individuals (described in more detail *infra*) or in *in vitro* cultures.

When mammalian cells are used as recombinant host cells for the production of polypeptides and RNA molecules, these cells will generally be grown in tissue culture. Methods for growing cells in culture are well known in the art (*see, e.g.*, Celis, J., ed., CELL BIOLOGY, Academic Press, 2nd edition, (1998); Sambrook, J. *et al.*, eds., MOLECULAR CLONING, A LABORATORY MANUAL, 2nd. edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989); Ausubel, F. *et al.*, eds., CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John H. Wiley & Sons, Inc. (1997); Freshney, R.,

-31-

CULTURE OF ANIMAL CELLS, Alan R. Liss, Inc. (1983)).

The selection of a host cell suited for a particular application will vary with a number of factors including the polypeptide or RNA molecule which is expressed. For example, when a glycoprotein is produced, it is generally desirable to express this protein in a cell type which will glycosylate the protein in a manner similar to that of the native protein.

In one aspect, the present invention provides methods for producing polypeptides and RNA molecules comprising introducing nucleic acid molecules of the invention into recombinant host cells and incubating these cells at a permissive temperature. In a related aspect, the invention provides purified polypeptides and RNA molecules produced according to the methods of the present invention.

Depending on the molecule which is expressed, the molecule may be obtained either from the culture supernatant or by lysing the recombinant host cells. When the expression product is a protein, it will often be possible to obtain the expression product from the culture supernatant. This will be so even when the protein does not have a naturally associated secretory signal. Codons encoding such a signal can be added to the vector sequences of the invention and will result in the expression of a fusion protein which will be secreted from the recombinant host cell. Nucleotide sequences encoding such leader sequences are known in the art and are publically available (*see, e.g.*, pPbac and pMbac vectors, STRATAGENE 1997/1998 CATALOG, Catalog #211503 and #211504, Stratagene, 11011 North Torrey Pines Road, La Jolla, CA 92037, USA).

Host cells may also be infected with packaged or unpackaged RNA molecules which have either been transcribed from the DNA molecules of the invention or replicated from such transcribed molecules. Further, these host cells may be infected at a restrictive temperature and then later shifted to a permissive one to activate expression of the gene of interest. The gene product of interest may then be recovered and purified by any suitable means.

The protein expressed from the gene of interest can be recovered and purified from recombinant cell cultures by methods known in the art including ammonium sulfate precipitation, anion or cation exchange chromatography.

-32-

phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography, and high performance liquid chromatography. Methods for purifying proteins are described in numerous sources (*see, e.g.*, Celis, J., ed., CELL BIOLOGY, Academic Press, 2nd edition. (1998)).

Methods for purifying RNA molecules are also known in the art (*see, e.g.*, Celis, J., ed., CELL BIOLOGY, Academic Press, 2nd edition. (1998)). These methods include phenol/chloroform extraction, digestion with DNAses followed by precipitation of the undigested RNA molecules, and column chromatography (*e.g.*, oligo dT column chromatography). Further, RNA molecules can be separated from other cellular material using the single-step guanidinium-thiocyanate-phenol-chloroform method described in Chomczynski and Sacchi, *Anal. Biochem.* 162:156-159 (1987).

A number of different bioprocess parameters can be varied in order to increase the amount of expression product produced during the cell culture process. The conditions under which the host cells are grown (*e.g.*, medium composition, pH, oxygen concentration, agitation, and, for the case of anchorage-dependent cells, the surface provided and the carrier of that surface) prior to exposure to the nucleic acid molecules of the invention or induction of gene expression influence both the cell density achieved at a given time and the physiological state of the cells. These culture conditions will thus affect the expected cellular response to vector exposure or the induction signal (*e.g.*, shifting to a permissive temperature). Further, the cell culture process-conditions mentioned above can be varied to maximize the production of expression product and, often, the characteristics (*e.g.*, glycosylation pattern) of that expression product.

The overall cell culture process employing nucleic acid molecules of the invention for the production of expression product can be implemented in a variety of bioreactor configurations (*e.g.*, stirred-tank, perfused, membrane enclosed, encapsulated cell, fluidized bed, and air-lift reactors) and scales (from laboratory T-flasks to thousands of liters), chosen to accommodate the requirements of the host cell line utilized (*e.g.*, anchorage dependency, O₂

-33-

concentrations). to maximize the production of expression product. and to facilitate subsequent recovery and purification of expression product.

The invention is also directed to the production of proteins or RNA molecules of interest using mammalian cells grown in serum-free or protein-free culture media. For example, by long-term culture under conditions restricting serum access or selecting for suspension growth, CHO cell lines are selected which are able to grow in serum-free medium and/or in suspension (Zang, M. *et al.*, *Bio/Technology* 13:389 (1995)). Further, by genetic modification of CHO K1 cells, a modified cell line designated CHO K1:cycE was obtained which grows as suspended single cells in protein-free culture media (Renner W. *et al.*, *Biotech. Bioeng.* 47:476 (1995)). CHO mutants (*e.g.*, LEC10 cells) have also been isolated which produce glycoproteins having different glycosylation patterns than those produced in parental CHO cells (Stanley, P., *Glycobiology* 2:99 (1992)). Alternatively, CHO cells capable of synthesize glycoproteins with correspondingly modified oligosaccharides may be obtained by genetically modifications which alter the activities of enzymes involved in oligosaccharide biosynthesis (Minch *et al.*, *Biotechnol. Prog.* 11:348 (1995)).

Further, a number of different bioprocess parameters can be varied in order to alter the glycosylation pattern of polypeptide products produced by the recombinant host cells of the invention. These factors include medium composition, pH, oxygen concentration, lack or presence of agitation, and, for the case of anchorage-dependent cells, the surface provided. Thus, the glycosylation pattern of glycoproteins may be altered by choosing the host cell in which these proteins are expressed in and the conditions under which the recombinant host cells are grown.

As explained below, polypeptides and RNA molecules of interest may also be produce in genetically engineered, non-human animals.

Gene Therapy

The vectors of the invention are also useful for gene therapy. When the vectors of the present invention are introduced into cells for gene therapy, the methods and vectors used will generally provide for the stable transfer of vector

-34-

sequences to the recombinant host cells. In such cases, vector sequences will be maintained in the host cell and will be transferred to cellular progeny. For example, the inclusion of long terminal repeats of retroviruses in gene transfer vectors has been found to confer stable maintenance of vectors sequences in recombinant host cells (Peng, L. *et al. J. Surg. Res.* 69:193-198 (1997); Qing, K. *et al., J. Virol.* 71:5663-5667 (1997)). Thus, chromosomal integration of vector sequences is one mechanism by which such sequences can be stably maintained in recombinant host cells. These sequence can integrate into host cell chromosomes either without regard to chromosomal location or at one or more specific chromosomal loci (*e.g.*, homologous recombination). These recombinant host cells may then be cultured *in vitro* or introduced into an individual.

The invention provides methods for expressing a sequence of interest in an individual to produce a polypeptide or RNA of interest comprising introducing nucleic acid molecules of the invention into host cells of the individual and regulating the temperature of the recombinant host cells. For example, vectors of the invention which express a "hot" sensitive replicase and contain a sequence encoding a polypeptide or RNA of interest can be introduced into human keratinocytes, epithelial cells, or fibroblasts *in vitro* and then reintroduced into a human subject. In such an instance, expression of the polypeptide or RNA of interest occurs when the temperature of tissues containing these cells is lowered to a permissive temperature.

The present invention also provides methods for administering polypeptides or RNA molecules to individuals in need thereof comprising introducing nucleic acid molecules of the invention into host cells, introducing the resulting recombinant host cells into these individuals, and inducing expression of the polypeptides or RNA molecules of interest. Similarly, host cells nucleic acid molecules of the invention can be introduced into host cells of an individual *in vivo*.

Induction of gene expression in individuals occurs by changing the temperature from a restrictive one to a permissive one. When the individual undergoing gene therapy is a human, and it is desirable for expression of the gene of interest to be activated only at specific times, 37°C will normally be a

-35-

restrictive temperature and gene induction will result from raising or lowering the temperature to a permissive one. In a similar fashion, when it is desirable for expression of the gene of interest to be inactivated only at specific times, 37°C will normally be a permissive temperature and gene inactivation will result from raising or lowering the temperature to a restrictive one.

The recombinant host cells introduced into an individual may be of any cell type that will at least be temporarily maintained in the individual or any cell type that will be maintained and at least temporarily retain and express nucleotide sequences of the invention. When the individual into which the recombinant host cells are introduced is a human, the host cells may be of any type which may be implanted in an area where the temperature may be altered between a permissive and a restrictive one by external means. For example, the recombinant host cells may be keratinocytes, epithelial cells, or fibroblasts which have been removed from an individual, transfected with a vector of the present invention, and reimplanted in an area near the surface where the skin temperature normally remains at or close to 37°C (e.g., an axilla). In such an instance, gene expression can be activated by altering the temperature of tissues containing the recombinant keratinocytes or fibroblasts to a permissive one (e.g., by placing an ice pack or peltier element over the location containing the recombinant host cells). Thus, the induction of expression of the gene of interest requires that the temperature of only a portion of the individual's body (e.g., axilla, arm, leg, hand, foot, neck region, etc.) be changed from a restrictive one to a permissive one.

Recombinant host cells may also be implanted in mammals at locations below surface, cutaneous tissues. One advantage to introducing recombinant host cells in such regions is derived from the temperatures of these tissues being more stably maintained than with surface, cutaneous tissues and, thus, gene expression is less likely to be activated by factors such as changes in climatic conditions. While the locations of suitable regions will vary with a number of factors, including the individual and the individual's normal body temperature, suitable tissues will generally include skin, nervous, and muscle tissues.

In another aspect, the invention provides methods for administering a polypeptide or RNA molecule to an individual in need thereof comprising the *in*

-36-

vivo introduction of nucleic acid molecules of the invention into host cells of the individual and inducing expression of heterologous polypeptides or RNA molecules encoded by these nucleic acid molecules. Methods for the *in vivo* introduction of alphaviral vectors to mammals tissues are described in Altman-Hamamdzc S. *et al.* (*Gene Ther.* 4:815-822 (1997)).

In a further aspect, methods are provided for administering a polypeptide or RNA molecule to an individual in need thereof by introducing RNA molecules to the cells of the individual. These RNA molecules may be obtained by a variety of methods including *in vitro* transcription and recombinant host cell expression. The RNA molecules may be introduced into cells of the individual either *in vitro* or *in vivo*. Methods for the introducing RNA sequences into host cells of individuals are described in Felgner, P. *et al.*, U.S. Patent No. 5,580,859.

The invention also provides non-infective, packaged RNA molecules encoding a temperature sensitive replicase useful as gene therapy vectors. These vectors have the advantages of being non-infectious, non-integrating, and express the gene of interest in a temperature-sensitive manner. Vectors of this type are useful for a variety of applications where a single administration of the gene product of interest is desired (*e.g.*, vaccine administration).

The nucleic acid molecules of the invention are useful for the regulated expression of stably integrated heterologous sequences in individuals. In one application, keratinocytes or fibroblasts of a human individual afflicted with diabetes are removed by tissue biopsy. DNA molecules of the invention containing a sequence of interest encoding human insulin are introduced and stably integrated into these cells *in vitro*. These recombinant host cells are reimplanted in a location near the surface where body temperature is relatively stably maintained (*e.g.*, an axilla). Prior to meal time, or some other time when insulin production is desired, the individual places an ice pack or a peltier element for a specified period of time over the location containing the recombinant host cells to induce expression of the heterologous insulin coding sequences. Further, a warm item may be used by the individual to raise the temperature to a permissive one when a cold sensitive replicase is used.

The actual temperature of the item which is placed in contact with the skin

-37-

will vary with the type of temperature-sensitive mutation used, the individual, the location of the recombinant host cells, the level of gene expression desired, and other factors.

Genetically Engineered, Non-Human Animals

5 Genetically engineered animals are currently used for the production of heterologous proteins (*see, e.g., Jeng, S. et al., J. Dairy Sci. 80:3167-3175 (1997); Limonta J. et al., Immunotechnology 1:107-113 (1995)*). These proteins are often harvested from bodily fluids such as blood, milk and urine (Meade, H. *et al., Nat. Biotechnol. 16:21-22 (1998); Kerr, D. et al., Nat. Biotechnol. 16:75-79 (1998)*).

10 The present invention also provides genetically engineered, non-human animal comprising cells which contain nucleic acid molecules of the present invention. These animals will generally have one or more DNA molecules of the invention stably integrated into their somatic and germ line cells. A number of methods are known in the art for producing animals having DNA molecules of the invention in their germ line cells (*see, e.g., Hew, C. et al., U.S. Patent No. 5,545,808; Jolicoeur, P., U.S. Patent No. 5,574,206; Mintz, B., U.S. Patent No. 5,550,316; Wagner, T. et al., U.S. Patent No. 4,873,191*). For example, DNA molecules can be introduced by microinjection into a fertilized, mammalian oocyte between the one-cell and eight-cell stage of embryological development. These oocytes are then implanted in a suitable female to produce founder animals which will stably transmit the heterologous transgene through the germ line to the next generation. Southern blot analysis is generally used to determine whether the genome of any particular individual carries the heterologous DNA sequence.

25 The genetically engineered animals may also contain nucleic acid molecules of the invention exclusively in somatic cells. Host cells containing these molecules may be implanted into the animal or nucleic acid molecules may be introduced into host cells of the animal *in vivo*.

30 Expression of the gene of interest in the cells of a genetically engineered animal may be induced by altering the body temperature of all or part of the animal from a restrictive one to a permissive one. Thus, the choice of the animal

-38-

used will vary with a number of factors, including the restrictive and permissive temperatures of the replicase employed, the normal body temperature of the animal to be genetically engineered, and the gene of interest. These animals may be either warm-blooded or cold-blooded. For example, Hew, C. *et al.* (U.S. Patent No. 5,545,808) describes the production of transgenic fish which express nucleotide sequences linked to an "anti-freeze" gene promoter. Expression of a sequence of interest in such an animal containing a nucleic acid molecule of the invention can be regulated by changing the water temperature the fish is kept in between restrictive and permissive temperatures.

When a warm-blooded animal contains a nucleic acid molecule of the present invention, the normal body temperature of the animal may be either a restrictive one or a permissive one. Further, in many instances expression of the gene of interest will either be induced or repressed in only a portion of the animal at any one time. For example, when the normal body temperature of a warm-blooded animal is a restrictive temperature and the temperature sensitive replicase is "hot" sensitive, the animal may be kept under conditions in which its extremities (*e.g.*, feet, arms legs, etc.) or surface tissues are lowered to a permissive one.

When a warm-blooded animal having cells which contain a nucleic acid molecule of the invention has a normal body temperature which is a permissive one, the gene of interest will generally be expressed in cells in internal regions of the animal. Such animals will be particularly useful for expressing the gene of interest in mammary gland and urothelial tissues. Kerr, D. *et al.* (*Nat. Biotechnol.* 16:75-79 (1998)), for example, describe the production of transgenic animals which express a foreign gene in the cells of their urothelium. These animals excrete the foreign gene product in their urine. Thus, the product of the gene of interest is readily collectable from such animals. Similarly, expression of the gene of interest in mammary gland tissues can result in the gene product being excreted into the animal's milk.

The present invention thus further provides genetically engineered, non-human animals which contain nucleic acid molecules of the invention in at least some of their cells. Also provided are genetically engineered, non-human

animals which contain DNA molecules of the invention stably integrated into the genome of some or all the animal's cells. The invention also provides methods for producing genetically engineered, non-human animals comprising introducing cells containing nucleic acid molecules of the invention into these animals, introducing nucleic acid molecules of the invention into the cells of these animals *in vivo*, or introducing DNA molecules of the invention into germ line cells to produce transgenic animals containing the sequence of interest in their somatic and germ line cells.

Pharmaceutical Compositions

The invention further provides pharmaceutical compositions, comprising polynucleotides of the invention in solution with a physiologically acceptable carrier and in a therapeutically effective amount. The administration of these pharmaceutical compositions may, for example, result in expression of a polypeptide in tissues of an animal which is immunogenic and intended to function as a vaccination. Similarly, the sequence of interest may encode polypeptides or RNA molecules required for the treatment of an active affliction. The administration of a pharmaceutical composition of the invention will thus be intended to have a therapeutic effect in these instances.

The nucleic acid molecules and recombinant host cells of the invention will normally be administered to an individual in a pharmacologically acceptable carrier. A composition is said to be "pharmacologically acceptable" if its administration can be tolerated by a recipient individual. Further, the composition of the invention will be administered in a "therapeutically effective amount" (*i.e.*, an amount that produces a desired physiological effect).

As would be understood by one of ordinary skill in the art, when the DNA molecules or recombinant host cells of the invention are administered to an individual, they may be in a composition which contains salts, buffers, adjuvants, or other substances which are desirable for improving the efficacy of the composition. Examples of materials suitable for use in preparing pharmaceutical compositions are provided in numerous sources including REMINGTON'S PHARMACEUTICAL SCIENCES (Osol, A. ed., Mack Publishing Co., (1980)).

The therapeutic compositions of the present invention can be administered

-40-

by various art known means but will normally be administered by injection, infusion or other suitable physical methods. The compositions may alternatively be administered intramuscularly, intravenously, or subcutaneously. Components of compositions for administration include sterile aqueous (*e.g.*, physiological saline) or non-aqueous solutions and suspensions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Carriers or occlusive dressings can be used to increase skin permeability and enhance antigen absorption.

When recombinant host cells are administered to an individual, the number of cells or nucleic acid molecules required to provide a therapeutically effective amount will vary with such factors as the individual's condition, the proteins or RNA molecules intended to be expressed, and the size of the individual.

Examples

The following enzymes and reagents were used in the experiments described in the examples which follow: Two polymerase, dNTPs and restriction enzymes were obtained from Boehringer Mannheim (9115 Hague Road, Indianapolis, IN 46250). T4 DNA ligase, fetal calf serum (FCS), bacto-tryptone and yeast extract was obtained from Gibco BRL (P.O. Box 68, Grand Island, NY, 14072, USA). *Bsp*120 I was obtained from MBI Fermentas, Inc. (300 Pearl St. Buffalo, NY, 14202, USA). XL-1 Blue competent cells were obtained from Stratagene (11011 North Torrey Pines Road, La Jolla, CA, 92037, USA). DNA purification kits and Taq polymerase were obtained from QIAGEN, Inc., (9259 Eton Avenue, Chatsworth, CA, 91311, USA). HP-1 medium was obtained from Cell Culture Technologies (Glattbrugg, Switzerland). All standard chemicals were obtained from Fluka (980 South 2nd St., Ronkonkoma, NY, 11779, USA), Sigma Chemical Co. (P.O. Box 14508, St. Louis, MO 63178, USA), Aldrich (1001 West St. Paul Ave. Milwaukee, WI, 53233, USA) and all cell culture materials were obtained from Becton Dickinson & Co. (1 Becton Drive, Franklin

Lakes, NJ. 07417. USA).

Example 1

Construction of the pCYTts vector system:

5 Manipulations and sequencing of DNA were carried out by standard procedures. The mutations in nsP2 were introduced by PCR using the following oligonucleotides:

oligo-nsP2 1: 5'-AACATTGAAATCGATATTACAGGGG (SEQ ID NO:2),

oligo-nsP2 2: 5'-CGGGTTATGGTCGACCGGGC (SEQ ID NO:3),

oligo-nsP2 3: 5'-GTGCCCTCCCCTGAGTTTAAACAATTCAGGGCCGA

10 ACGCG (SEQ ID NO:4), and

oligo-nsP2 4: 5'-GAATTGTTTAAACTCAGGAGGCACCCTCGTGG (SEQ ID

NO:5), the single restriction sites used for first analysis and subsequent cloning (*DraI*, *Clal* and *Sall*) are underlined. PCR reactions were performed using either

15 oligo-nsP2 1 (SEQ ID NO:2) and oligo-nsP2 3 (SEQ ID NO:4) or oligo-nsP2 2 (SEQ ID NO:3) and oligo-nsP2 4 (SEQ ID NO:5). 100 pmol of each oligo was

used and 5 ng of the template DNA (pSinRep5: Xiong, C. *et al.*, *Science* 243:1188-1191 (1989)) was used in the 100 μ l reaction mixture, containing 4

units of Taq or Pwo polymerase, 0.1 mM dNTPs and 1.5 mM MgSO₄. All DNA concentrations were determined photometrically using the GeneQuant apparatus

20 (Pharmacia Biotech Inc., 800 Centennial Ave., Piscataway, NJ. 08854). The

polymerase was added directly before starting the PCR reaction (starting point was 95°C). The temperature cycles were as follows: 95°C for 2 minutes,

followed by 5 cycles of 95°C (45 seconds), 58°C (30 seconds), 72°C (90 seconds) and followed by 25 cycles of 95°C (45 seconds), 68°C (30 seconds),

25 72°C (90 seconds).

The two PCR fragment were purified using Qia spin PCR kit (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA. 91311) and finally digested in an appropriate buffer using 20 units of *Sall* and *DraI*, respectively 20 units of *Clal* and *DraI*. The digestion was performed for 12 hours at 37°C. The DNA

-42-

fragments were gel-purified (Gene-Clean; Bio 101 Inc., 1070 Joshua Way, Vista, CA, 92083, USA) and finally ligated into *Clal/SaII* digested and gel-purified SinRep5 vector (Xiong, C. *et al.*, *Science* 243:1188-1191 (1989). The correct sequence of the obtained vector was checked by DNA sequencing of the whole nsP2 gene.

The mutations in nsP4 were also introduced by PCR using the following oligonucleotides:

oligo-nsP4 1: 5'-GGTAGACGAGACAGTCGCATGCCTGGATAC (SEQ ID NO:6),

oligo-nsP4 2: 5'-GTATCCAGGCATGCGACTGTCTCGTCTACC (SEQ ID NO:7),

oligo-nsP4 3: 5'-CAGACCGGTTAACGCCATAGCGTCG (SEQ ID NO:8), and

oligo-nsP4 4: 5'-CTCTATTACTAGTATGGACAGTTGG (SEQ ID NO:9), the

singular restriction sites used for the first analysis and the final cloning step (*SphI*,

HpaI and *SpeI*) are underlined. Two PCR reactions were carried out as described

above using either oligo-nsP4 1 (SEQ ID NO:6) and oligo-nsP4 3 (SEQ ID NO:8)

or oligo-nsP4 2 (SEQ ID NO:7) and oligo-nsP4 4 (SEQ ID NO:9).

Both PCR products were gel-purified and then used in assembly PCR to amplify the whole nsP4 gene. For the assembly PCR, 50 pmol of the outer primers (3 and 4) and about 10 ng of each PCR fragment was used. The reaction volume was 100 μ l, containing 4 units of Taq or Pwo polymerase, 0.1 mM dNTPs and 1.5 mM MgSO₄. The PCR conditions were as followed:

Ninety-five °C for 2 minutes, followed by 5 cycles of 92 °C (45 seconds), 58 °C (30 seconds), 72 °C (120 seconds) and followed by 25 cycles of 92 °C (45 seconds), 64 °C (30 seconds), 72 °C (120 seconds).

The obtained PCR fragment was purified as described above and the eluate was digested with 20 units of *SpeI* and *HpaI* in an appropriate buffer. The fragment was gel-purified and ligated into gel-purified *SpeI/HpaI* restricted SinRep5 vector. The correct sequence of the obtained vector was checked by DNA sequencing.

Over night digestion of SinRep5-nsP4mut and SinRep5-nsP2mut with *SpeI/HpaI* and gel purification of the nsP4 fragment and sinRep-nsP2mut vector.

-43-

The nsp4mut fragment was ligated into the SinRep5-nsp2mut vector. The final step was cloning the nsp gene into the 987/SinRep5 vector (Bredenbeek, P. *et al.*, *J. Virol.* 67:6439-6446 (1993)) using *Clal* and *HpaI* as restriction endonucleases. the resulting vector was named pCYTts (FIG. 2 and FIG. 3A-3D (SEQ ID NO:1)).

pCYTts constructs: Five different genes were cloned into the pCYTts vector. Green fluorescent protein (GFP), secreted alkaline phosphatase (SEAP), β -interferon (β -INF), erythropoietin (EPO), and HIV gp160.

Example 2

Regulated expression of GFP in transient and stable expression

The pCYTts system was successfully used to express cytoplasmic proteins. as an example we used the green fluorescent protein (GFP) of the jellyfish *Aequorea victoria* (Cramer *et al.*, *Nat. Biotech.* 14:315-319 (1996)). GFP is ligated into pCYTts via *XbaI* and *Bsp120I* (Fermentas). Clones with the correct insert were identified by restriction digest. The GFP expression was tested in both. transient and stable expression.

Transient transfection in BHK 21 cells was carried out using the CaPO₄ precipitation transfection protocol: 6 μ g of plasmid DNA (pCYTts GFP) in 30 μ l H₂O was mixed with 30 μ l of an 1 M CaCl₂ solution. After addition of 60 μ l phosphate buffer (50 mM HEPES, 280 mM NaCl, 1.5 mM Na₂HPO₄, pH 7.05) the solution was vortexed for 5 seconds. followed by an incubation at room temperature for 25 seconds. The solution was immediately added to 2 ml HP-1 medium containing 2% FCS (2% FCS medium). The medium of a 80% confluent BHK21 cell culture in a 6-well plate was then replaced by the DNA containing medium. After an incubation for 5 hours at 37°C in a CO₂ incubator. the DNA containing medium was replaced by 2 ml of 15% glycerol in 2% FCS medium. The glycerol containing medium was removed after a 30 second incubation phase and the cells were washed with 5 ml of HP-1 medium containing 10% FCS. Finally 2 ml of fresh HP-1 medium containing 10% FCS

-44-

was added.

After transient transfection of BHK cells with pCYTsGFP, the expression was tested at 37°C. No expression of GFP was detected using the methods described below. GFP was produced when the temperature was shifted down to 29°C. The GFP expressing cells survived for at least 5 days.

Stable transfection in BHK21 cells. The stable transfection is carried out essentially as described for the transient transfection, except, for the stable transfection linearized plasmid DNA was used. Twenty μg of pCYTsGFP was incubated with 30 units of *NaeI* in an appropriate buffer for at least 4 hours at 37°C. The reaction was stopped by phenol/chloroform extraction, followed by an isopropanol precipitation of the linearized DNA. The restriction reaction was checked by gel electrophoresis using a 0.8% agarose gel, stained with ethidium bromide. For the transfection 5.4 μg of linearized pCYTsGFP was mixed with 0.6 μg of circular pSVtrpB (selection plasmid) in 30 μl H_2O . Followed by the procedure described for transient transfection.

Stably transfected cells were selected and grown in selection medium (HP-1 medium, without tryptophane, supplemented with 300 μM indole and 5% dialyzed FCS) at 37°C in a CO_2 incubator. When the mixed population had grown to confluency, the culture was divided into two parts and both parts were cultured for an additional 12 hours at 37°C. One part of the cells was then shifted to 30°C to induce the expression of the gene of interest. The other part was kept at 37°C.

Detection of gene expression

Green fluorescent protein can be easily detected in a spectrofluorometer, due to its strong fluorescence. This is seen when GFP is located in the cytoplasm of the cell. GFP production was detected by fluorescence microscopy and quantified by whole cell spectrofluorophotometry (Spectrofluorophotometer, Shimadzu RF-5001PC). Detached cells were washed with 5 ml PBS (per liter: 0.132 g $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$; 0.20 g KCl; 0.20 g KH_2PO_4 ; 0.10 g $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$; 8 g NaCl; 1.15 g Na_2HPO_4 ; pH 7.2) and resuspended in 1 ml PBS. The excitation

-45-

wavelength was 397 nm and the emission wavelength was 510 nm. To carry out the measurements in a linear range for fluorescence detection, the cells were diluted to obtain a fluorescence between 0.05 and 1.0 emission units.

To determine the optimal induction temperature, cultures of mixed populations of stable transfected cells were incubated for 48 hours at different temperatures in selection medium without FCS. Expression was induced when cultures were shifted to 34°C or lower. The highest expression was detected at 29°C (FIG. 4A). When stable transfected cells were induced at 30°C for 4 hours and subsequently grown at 37°C for 24 hours, green cells could be observed by fluorescence microscopy. This clearly showed that the expression of the gene of interest starts after 4 hours of induction (FIG. 6A).

Time dependence

The kinetics of the system were determined by photometrically at different time points after induction. GFP expression was detected as described above. Ten hours after induction a clear expression of GFP is detectable at 29°C (FIG. 5A). When shifting the cells back to 37°C after induction, new mRNA production should be blocked, however, the translation of the protein of interest should occur with a higher expression level. The cells were shifted after 4, 6, 8 or 10 hours after induction back to 37°C. 24 hours later the expression of GFP was detected as described above (FIG. 6A). Thus, transcription starts shortly after induction.

Long term stability of the cell line

To determine the long term expression of the gene of interest, stably transfected cells were cultured for at least 8 weeks at 37°C. The expression of GFP was tested by shifting the cells to 29°C. No difference was observed in the expression level of GFP between cells used directly after stable transfection and cells cultured for at least 4 weeks.

Example 3

Regulated expression of SEAP in transient and stable expression

The pCYTts system was successfully used to express secreted proteins, as an example we used the secreted alkaline phosphatase (SEAP) of human origin (CLONTECH Laboratories, Inc., 1020 East Meadow Circle, Palo Alto, CA, 94303, USA). The SEAP coding sequence is ligated into pCYTts via *Xba*I and *Stu*I. Clones with the correct insert were identified by restriction digest. SEAP expression was tested for both transient and stable expression.

Transient transfection in BHK21 cells was carried out using the CaPO₄ co-precipitation transfection protocol as described in Example 2.

Stable transfection in BHK21 cells

The stable transfection was carried out essentially as described for transient transfection, except that linearized plasmid DNA was used. Twenty μ g of pCYTtsSEAP was incubated with 30 units of *Mlu*I in an appropriate buffer for at least 4 hours at 37°C. 10 μ g of pSVneo was digested with 30 units of *Sca*I for at least 4 hours at 37°C. Both reactions were stopped by phenol/chloroform extraction, followed by an isopropanol precipitation of the linearized DNA. The restriction reactions were checked by gel electrophoresis using a 0.8 % agarose gel, stained with ethidium bromide. For the transfection 5.88 μ g of linearized pCYTtsSEAP is mixed with 0.12 μ g of linearized pSVneo (selection plasmid) in 30 μ l H₂O. Followed by the procedure described for the transient transfection.

Detection of gene expression

Transient and stable transfected cells containing pCYTtsSEAP were tested for SEAP expression after 3 days of induction by dot blotting. 2.5 μ l of cell culture supernatant was spotted on a nitrocellulose membrane. After drying the membrane for 10 minutes at room temperature, the development reaction was

-47-

carried out using alkaline phosphatase detection reagents (10 ml AP buffer (100 mM Tris/HCl, 100 mM NaCl, pH 9.5) with 50 μ l NBT solution (7.7% Nitro Blue Tetrazolium (Sigma) in 70% dimethylformamide) and 37 μ l of X-Phosphate solution (5 % of 5-bromo-4-chloro-3-indolyl phosphate in dimethylformamide).

5 The SEAP activity was quantified in an colorimetric enzymatic activity test. 500 μ l of culture supernatant containing SEAP was incubated at 65°C for 5 minutes, and finally centrifuged (20,000g; 20 seconds). To determine the SEAP activity 400 μ l of the centrifuged supernatant were mixed with 500 μ l of 2 x SEAP buffer (20 mM L-homoarginine, 2 M diethanolamine, and 1 mM
10 MgCl₂•6H₂O, pH 9.8) in a cuvette. The SEAP activity was followed in a spectrophotometer at 405 nm. after adding 100 μ l nitrophenylphosphate (120 mM) (Sigma 104, Sigma) to the sample. The absorbance was measured every 30 second over a time period of 10 minutes. The obtained values at different time points were plotted versus the time and a plot with a linear slope was obtained.

15 In the mixed population the amount of SEAP molecules produced per cell was estimated to be around 10⁷ molecules per cell. To get a stable expression of SEAP, cloned cells were automatically sorted in a cell sorter and finally analyzed for SEAP activity. About one out of 20 clones showed SEAP expression. The SEAP expression was estimated to be one order of magnitude higher than in the
20 mixed population.

 Highest SEAP expression was detected at 29°C (FIG. 4B). SEAP activity could be detected 15 hours after induction at 29°C (FIG. 5B). However, expression of SEAP started much earlier, as shown in FIG 6B. The SEAP expressing cells were shifted after 4, 6, 8 or 10 hours of induction back to 37°C,
25 24 hours later the expression of SEAP was detected as described above. SEAP expression could be detected as early as 6 hours after induction (FIG. 6B). Thus the SEAP expression also started shortly after induction.

Example 4

Regulated expression of β -INF in transient and stable expression

A β -interferon gene of human origin was used to demonstrate that the pCYTts system can be used to express antiviral, secreted proteins. β -interferon has antiviral activity and interferes with RNA replication. pCYTts systems tightly regulate the expression of genes even when these genes encode proteins which interfere with RNA replication.

The gene encoding β -interferon was generated as described in Prodrumou, C. and Pearl, L. (*Protein Eng.* 5:827-829 (1992)). Primers were generated using the human β -interferon nucleotide sequences disclosed in GenBank reports V00534, J00218, K00616, and M11029. The β -interferon cDNA was ligated into pCYTts after restriction with *Xba*I and *Bsp*120I. Expression of β -interferon was tested in transient and stable expression systems.

Transient and stable (mixed population) expression of β -INF was determined by Western-blotting. 0.5 ml of culture medium was methanol/chloroform precipitated and the pellet was resuspended in SDS-PAGE sample buffer. Samples were heated for 5 minutes at 95°C before being applied to 15% acrylamide gels. After SDS-PAGE, proteins were transferred to Protan nitrocellulose membranes (Schleicher & Schuell, Inc., 10 Optical Ave., Keene, NH 03431, USA). The membrane was blocked with 1 % bovine albumin (Sigma) in TBS (10xTBS per liter: 87.7 g NaCl, 66.1 g Trizma hydrochloride (Sigma) and 9.7 g Trizma base (Sigma), pH 7.4) for 1 hour at room temperature, followed by an incubation with a mouse anti-human β -INF antibody (0.2 μ g/ml, Research Diagnostics Inc., USA) for 1 hour. The blot was washed 3 times for 10 minutes with TBS containing 0.05% Tween20 (TBS-T), and incubated for 1 hour with a horseradish peroxidase-anti-mouse IgG conjugate (0.1 μ g/ml, Amersham Life Science, England). After washing 2 times for 10 minutes with TBS-T and 2 times for 10 minutes with TBS, the development was carried using the ECL kit (Amersham).

Samples for the blot were taken after 3 or 5 days incubation at 29°C.

-49-

Another part of the culture was kept at 37°C and a sample was taken after 5 days. FIG. 8A shows that β -INF is produced after 3 days at 29°C whereas incubation at 37°C yields no detectable β -INF production.

Example 5

Regulated expression of EPO in transient expression

The pCYTts system was successfully used to express pharmaceutically relevant, secreted proteins. As an example of such expression, we used a gene of human origin encoding erythropoietin (EPO). This gene was generated by PCR as described in Example 4. Primers were generated using the human EPO nucleotide sequences disclosed in GenBank report X02158. The gene encoding EPO was ligated into pCYTts following restriction with *Xba*I and *Bsp*120L (Fermentas). Clones with the correct insert were identified by restriction digest. EPO expression was tested in both transient and stable expression systems.

BHK21 cells were transiently transfected according to the CaPO_4 co-precipitation protocol, as described in Example 2.

EPO production was determined by western blotting, as described in Example 4. The detection was carried out by incubating the nitrocellulose membrane with 2 μg rabbit anti-human EPO antibody (Research Diagnostics Inc.) in 10 ml TBS-T for 1 hour, followed by 3 washes, each for 10 minutes, with TBS-T. Finally, the nitrocellulose membrane was incubated for 1 hour with alkaline phosphatase conjugated anti-rabbit IgG (Jackson ImmunoResearch Laboratories, Inc.) diluted 1:5000 in TBS-T. After washing 2 times for 10 minutes with TBS-T and 2 times for 10 minutes with TBS, the blot was developed by alkaline phosphatase staining as described in Example 3. Transiently transfected cells induced for 4 days at 29°C produced detectable amounts of EPO (FIG. 8B).

-50-

Example 6***Regulated expression of EPO in stable expression***

In the pCYTs504EPO expression vector, the human erythropoietin (EPO) coding sequence (including its natural leader peptide for secretion into the growth medium) was fused in frame to the sequence coding for the Sindbis virus capsid protein (C-protein). The rationale of this construct was to include the translational enhancer located within the C-protein coding region that has been shown to lead to a 10- to 20-fold increased expression level compared to constructs lacking this enhancer (Frolov *et al.*, *Proc. Natl. Acad. Sci. USA* 93:11371-11377 (1996)). The fusion gene is expressed from the subgenomic promoter of expression vector pCYTs. Upon co-translational release of the EPO precursor from the fusion protein, catalyzed by the autoproteolytic activity of the C-protein, EPO is directed to the secretory pathway by its N-terminal leader peptide.

For stable transfection, a 3:1 ratio of pCYTs504EPO (linearized by restriction cleavage with *Mlu*I) and the neomycin resistance-conferring plasmid 987BBneo (Bredenbeek *et al.*, *J. Virol.* 67:6439-6446 (1993) (linearized by restriction cleavage with *Sca*I) were introduced into BHK21 cells using the calcium phosphate co-precipitation protocol described in Example 2. After 1 week incubation at 37°C under selective conditions (HP-1 medium supplemented with 10% FCS and 200 µg/ml G418 (neomycin)), single colonies were separated and further propagated under the same conditions.

To screen for EPO-secreting clones, cells were grown in 12-well plates at 37°C to 80% confluency and incubated at 30°C for further 4 days. Three µl of each culture supernatant were analyzed for secreted EPO by Dot Blot analysis using an anti-EPO rabbit IgG and an anti rabbit IgG-alkaline phosphatase conjugate. Among 27 clones investigated, one EPO-secreting clone was identified. A rough concentration of 2.5 mg EPO per liter of supernatant was estimated using an EPO ELISA Kit (Boehringer Mannheim). The identity of the secreted protein was further confirmed by Western Blot analysis. For that

-51-

purpose, cells were grown to 80% confluency at 37°C in a T-75 cell culture flask with 30 µl HP-1 medium (without FCS) supplemented with G418 (200 µg/ml) and then incubated at 30°C for further 4 days. Twenty µl of the culture supernatant were separated on a 15% SDS polyacrylamide gel and blotted onto a nitrocellulose membrane. Using an anti-EPO rabbit IgG/anti-rabbit IgG-alkaline phosphatase conjugate system, a single protein was specifically detected, that showed the same electrophoretic mobility as an authentic EPO sample from a different source (apparent M_r about 29 kDa) (FIG. 9). The resulting cell line was named 1C4.

Example 7

Production of Sindbis Virus particles containing EPO RNA

One µg of RNase-free vector (pDH-EB; Bredenbeek *et al.*, *J. Virol.* 11:6439-6446 (1993)) was linearized by *EcoRI* digestion. Subsequently *in vitro* transcription was carried out using the SP6 *in vitro* transcription kit (InvitroscriptCAP by Invitrogen, Invitrogen BV, NV Leek, Netherlands). The resulting 5'-capped mRNA was analyzed on reducing agarose-gels.

Five µg of *in vitro* transcribed mRNA were electroporated into 1C4 cell line (Example 6) according to Invitrogen's manual (Sindbis Expression system, Invitrogen BV, Netherlands). After 10 hours incubation at 37°C the FCS containing medium was exchanged by HP-1 medium without FCS, followed by an additional incubation at 30°C for 72 hours. The supernatant was passaged to a BHK 21 cell layer, incubated for 2 hours at 4°C and finally discharged. The cells were washed 4 times with HP-1 buffer and incubated for 24 hours at 30°C. Three µl of the culture supernatant were analyzed for secreted EPO by Dot Blot analysis using an anti-EPO rabbit IgG and an anti rabbit IgG-alkaline phosphatase conjugate (FIG. 10).

-52-

Example 8***Regulated expression of gp160 in transient expression***

The pCYTts system was used to express gp160, the HIV envelope protein. The gp160 gene was amplified from pAbT4674 (ATCC 40829) and cloned in pCYTts via *Xba*I and *Bsp*120L. BHK21 cells were transiently transfected using lipofectamine (Life Technologies, Basel, Switzerland). 0.8 µg pCYTts gp160 DNA in 150 µl Dulbecco's Modified Eagle medium (DMEM, Life Technologies, Basel, Switzerland) was mixed with 150 µl DMEM containing 2.5 µl lipofectamine. The solution was incubated at room temperature for 15 minutes and added to a 80% confluent BHK cell layer in a 24-well plate. After incubation for 5 hours at 37°C in a CO₂ incubator, the cells were washed and incubated for another 12 hours at 37°C. Cells were split and one part was incubated at 29°C and one part was incubated at 37°C for 5 days. Cells were harvested and lysed in SDS-PAGE sample buffer. Samples were heated for 5 minutes at 95°C and applied to a 8% acrylamide gel. gp160 expression was analyzed by Western blotting as described in Example 4. The nitrocellulose membrane was incubated with rabbit anti-human gp160 antibody (kindly provided by Dr. Schawaller, Diamed AG, Switzerland), diluted 1:3000 in 10 ml TBS-T for 1 hour and subsequently washed three times for 10 minutes with TBS-T. Then the membrane was incubated for 1 hour with alkaline phosphatase conjugated goat anti-rabbit IgG (Jackson ImmunoResearch Laboratories, Inc.) diluted 1:5000 in 10 ml TBS-T. The membrane was washed two times with TBS-T for 10 minutes and two times for 10 minutes with TBS. Development was carried out as described in Example 3. Transiently transfected cells produced detectable amounts of gp160 (data not shown).

Example 9

Regulated expression of GFP in human foreskin fibroblasts

The pCYTts system was used to express green fluorescent protein in human foreskin fibroblasts. Cells were transfected using lipofectamine as described in Example 8. Twelve hours post-transfection one part of the cells was incubated at 37°C, the other part was incubated at 29°C. After 48 hours bright green cells were observed by fluorescence microscopy in the cultures incubated at 29°C, whereas at 37°C no GFP expression was detected.

Example 10

A multivector system with the insert in sense direction

The regulatable vector system of the invention was used for the production of non-cytopathic viral particles. As the gene of interest we chose the structural proteins of Sindbis virus and as a marker protein we chose GFP. The cells were stably transfected with pCYTtsGFP, as described in Example 2. The stable transfected cells were transiently transfected with a defective helper construct (pDHBB; Bredenbeek *et al.*, *J. Virol.* 11:6439-6446 (1993)), carrying the Sindbis virus structural proteins according to the protocol described in Example 2.

Transfected cells were grown overnight at 37°C. The cells were then shifted to 29°C to induce viral gene expression. The viral particles formed contain packaged pCYTtsGFP RNA sequences, and GFP is expressed when the packaged viral particles infect new target cells. To perform the new infection, the medium was collected and centrifuged (1800 rpm; 3 minutes) after 4 days of expression. The supernatant was placed on 80% confluent BHK cell layers and incubated for 4 hours at 29°C. After the incubation phase the medium was discharged and the cells were washed 3 times with 5 ml HP-1 medium followed by incubation at 29°C for an additional 24 hours. Finally, the expression level

-54-

of the marker gene GFP was measured by fluorescence spectroscopy, as described in Example 2.

About 10% of the BHK cells initially produced GFP and after an additional 24 hours of incubation at 29°C all cells of the expressed GFP. The conditioned medium of these cells was again harvested, centrifuged and placed onto a 80% confluent layer of BHK cells. After 48 hours of incubation at 29°C, 100% of these cells were found to express GFP.

As a control, the transfected cells were grown for 5 days at 37°C after which conditioned medium was collected and centrifuged (1800 rpm, 3 minutes). The supernatant was placed onto an 80% confluent BHK cell layer. After 8 hours of incubation at 29°C, the medium removed and the cells were washed 3 times with 5 ml HP-1 medium and incubated at 29°C for additional 24 hours. Finally, the expression level of GFP is determined. No GFP expressing cells could be detected (FIGs. 7A and 7B).

Example 11

A multivector system with insert in antisense direction

As a model system we tested the regulatable system with the production of viral particles. As the gene of interest we choose the structural proteins of Sindbis virus and as a marker protein we chose GFP. The cells were stably transfected with pCYTsGFP, as described in Example 2.

The antisense helper vector was constructed as follows:

The structural proteins were obtained by digesting the pDHBB vector (Bredenbeck, P. *et al.*, *J. Virol.* 67:6439-6446 (1993)) with *EcoRI* and *BamHI*. The fragment was purified by gel electrophoresis and cloned into *EcoRI/BamHI* digested pMPSVHE vector (Artelt, P. *et al.*, *Gene* 68:213-219 (1988)). Since the *EcoRI* and *BamHI* restriction sites are in opposite orientations in these vectors, the structural protein fragment was cloned in an antisense orientation into pMPSVHE. The resulting vector was named pMPSVanti-DHBB.

The stable transfected cells were transiently transfected with

-55-

pMPSVanti-DHBB as described in Example 10. Transfected cells were grown overnight at 37°C. The cells were then shifted to 29°C to induce viral gene expression. After 4 days of induction, the conditioned medium was collected and centrifuged (1800 rpm; 3 minutes). The supernatant was placed on 80% confluent BHK cell layers and was incubated for 4 hours at 29°C. After the incubation phase the medium was discharged and the cells were washed 3 times with 5 ml HP-1 medium and incubated at 29°C for additional 24 hours. Finally, the expression level of the marker gene GFP was measured by fluorescence spectroscopy, as described in Example 2.

About 1% of the BHK cells initially produced GFP and after an additional 120 hours incubation at 29°C 30% of the cells express GFP. Thus, even antisense DNA fragments can be used within this invention to produce functional proteins.

Conclusions

The expression system described in the preceding examples fulfills nearly all of the criteria for an ideal inducible gene expression system as described in Saez, E. *et al.*, (*Curr. Opin. Biotechnol.* 8:608-616 (1997)). This system is very specific in that it is only switched on when the temperature is shifted to below 34°C. The basal expression, as shown in several experiments, is not detectable with the standard detection methods used in the preceding examples. Even with the very sensitive system of viral infection (FIG. 7A-7B and FIG. 9) no basal expression at 37°C could be detected. This shows the high degree of regulatory stringency, because a functional replicase molecule would initiate an autocatalytic cycle of RNA replication and transcription which would result in a high expression level of the protein of interest.

Further, as shown in FIG. 6A-6B, gene expression starts rapidly after induction and stops quickly after the temperature is shifted back to a restrictive one. There is no problem with the bioavailability of the inducer, because temperature shifts to 29°C rapidly disseminate and are non-toxic. Once a restrictive temperature has been reached, the duration of gene expression is only

-56-

dependent on the stability of the mRNA encoding the protein or RNA of interest.

Compared with the tetracycline system, the system described in the preceding examples has the advantages that there is no detectable basal expression and the bioavailability of a temperature shift is much less harmful than the antibiotic tetracycline or the expression of the tTA protein. A further advantage of the herein described regulatable DNA vector system is that only one vector need be introduced into a host cell, because all relevant proteins needed for the expression and regulation can be encoded by this one vector. This is in contrast with the tetracycline system where two vectors must be transfected into the cells (Gossen, M. & Bujard, H., *Proc. Natl. Acad. Sci. USA* 89:5547-5551 ((1992))).

As already noted, the turning off of the pCYTs system is dependent on the stability of the replicase and the mRNA encoding the protein of interest. It has been shown that the half-life of the replicase is one half hour after expression (De Groot *et al.*, *Proc. Natl. Acad. Sci. USA* 88:8967-8971 (1991)). The mRNA stability is therefore the limiting factor which determines how rapidly the system is switched off. SEAP mRNA, for example, was translated for about 10 hours after shifting to restrictive temperature (FIG. 6A-6B). This high stability has also been found for CAT mRNA (Xiong, C. *et al.*, *Science* 243:1188-1191 (1989)), suggesting that mRNA derived from the Sindbis virus is very stable regardless of the protein encoded by this mRNA.

The system was tested in mixed population to prove that the expression system is independent of the site of integration and the copy number, as shown in FIG. 4A-4B and FIG. 5A-5B.

In conclusion, the pCYTs temperature regulatable gene expression system described in the preceding examples has significant advantages over the commonly used regulatable systems. Due to its very low level of basal expression, this system can be used for the expression of host toxic proteins, as shown with the expression of the HIV envelope protein gp160, which so far has been a difficult task with previous *in vitro* gene expression systems. Since the present system has also tested for the long term expression and the reinducibility, it is useful for gene therapy. Its potential use for gene therapy has been shown

-57-

in the transient expression of GFP in human skin cells, which are easy accessible for temperature regulation.

It will be clear that the invention may be practiced otherwise than as particularly described in the foregoing description and examples.

5 Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, are within the scope of the appended claims.

10 The entire disclosure of all publications (including patents, patent applications, journal articles, laboratory manuals, books, or other documents) cited herein are hereby incorporated by reference.

-58-

What Is Claimed Is:

1. A DNA molecule comprising a polynucleotide which encodes an RNA molecule, said RNA molecule comprising:

(a) at least one *cis*-acting sequence element,

5 (b) a first open reading frame having a nucleotide sequence encoding a non-cytopathic, temperature-sensitive RNA-dependent RNA polymerase, and

(c) at least one second nucleotide sequence selected from the group consisting of:

10 (i) a second open reading frame encoding a protein, or portion thereof, wherein said second open reading frame is in a translatable format after one or more RNA-dependent RNA replication events;

(ii) a sequence complementary to all or part of the second open reading frame of (i); and

15 (iii) a sequence encoding an untranslated RNA molecule, or complement thereof.

2. A DNA molecule of claim 1, wherein the RNA-dependent RNA polymerase is of viral origin.

3. A DNA molecule of claim 1, wherein the RNA-dependent RNA
20 polymerase is of alphaviral origin.

4. A DNA molecule of claim 1 which encodes an RNA-dependent RNA polymerase that has replicase activity at temperatures below 34°C and low or undetectable replicase activity at 34°C and above.

5. A DNA molecule of claim 1, wherein the second open reading
25 frame of 1(c)(i) encodes a cytokine, lymphokine, tumor necrosis factor, interferon, toxic protein, prodrug converting enzyme, or other protein.

-59-

6. A DNA molecule of claim 1, wherein the second open reading frame of 1(c)(i) encodes human erythropoietin or human β -interferon.

7. A DNA molecule of claim 1, wherein the second nucleotide sequence of 1(c)(ii) or 1(c)(iii) encodes an untranslated RNA molecule selected from the group consisting of an antisense RNA molecule, tRNA molecule, rRNA molecule, ribozyme.

8. A method of making a recombinant host cell comprising introducing a DNA molecule of claim 1 into a host cell.

9. An *in vitro* cell culture comprising a recombinant host cell produced by the method of claim 8.

10. An *in vitro* cell culture comprising a recombinant host cell comprising a DNA molecule of claim 1.

11. The cell culture of claim 10, wherein some or all of the DNA sequences of a DNA molecule of claim 1 are stably maintained in said host cell.

12. An RNA molecule transcribed from a DNA molecule of claim 1.

13. An alphaviral particle containing an RNA molecule of claim 12.

14. An *in vitro* cell culture comprising a recombinant host cell comprising an RNA molecule of claim 12.

15. An isolated nucleic acid molecule comprising a polynucleotide having the nucleotide sequence of SEQ ID NO:1.

-60-

16. A method for producing a protein or untranslated RNA molecule in a recombinant host cell comprising:

- (a) growing host cells under suitable culture conditions;
- (b) introducing at least one DNA molecule of claim 1 into said
- 5 host cells; and
- (c) recovering said protein or untranslated RNA molecule.

17. A method for producing a protein or untranslated RNA molecule in a recombinant host cell comprising:

- (a) growing host cells under suitable culture conditions;
- 10 (b) introducing at least one RNA molecule of claim 12 into said host cells; and
- (c) recovering said protein or untranslated RNA molecule.

18. The method of claim 17, wherein the protein is erythropoietin.

19. The method of claim 17, wherein said RNA is packaged into an

15 alphaviral particle.

20. A method for producing alphaviral particles containing an RNA molecule of claim 12 comprising:

- (a) growing host cells under suitable culture conditions;
- (b) introducing into said host cells at least one DNA molecule
- 20 of claim 1 having one or more open reading frames which encode alphaviral structural proteins; and
- (c) recovering said alphaviral particles.

21. A method for producing a protein in a recombinant host cell comprising:

- 25 (a) growing host cells under suitable culture conditions;
- (b) infecting said host cells with alphaviral particles produced

-61-

by the method of claim 20: and

- (c) recovering said protein.

22. The method of claim 21, wherein said protein is erythropoietin.

23. A method for regulating the expression of a protein or
untranslated RNA molecule in a recombinant host cell comprising:

- (a) growing host cells under suitable culture conditions;
(b) introducing at least one DNA molecule of claim 1 into said
host cells; and
(c) changing the temperature of the host cell culture from:
(i) a permissive temperature to a restrictive
temperature, or
(ii) a restrictive temperature to a permissive
temperature.

24. A method for regulating the expression of a protein or
untranslated RNA molecule in a recombinant host cell comprising:

- (a) growing host cells under suitable culture conditions;
(b) introducing at least one RNA molecule of claim 12 into
said host cells; and
(c) changing the temperature of the host cell culture from:
(i) a permissive temperature to a restrictive
temperature, or
(ii) a restrictive temperature to a permissive
temperature.

25. A method for regulating the expression of a protein or
untranslated RNA molecule in an individual comprising:

- (a) administering at least one DNA molecule of claim 1 to
said individual: and
(b) changing the temperature of at least a portion of said

-62-

individual from:

- (i) a permissive temperature to a restrictive temperature, or
- (ii) a restrictive temperature to a permissive temperature.

26. A method for regulating the expression of a protein or untranslated RNA molecule in an individual comprising:

- (a) administering at least one RNA molecule of claim 12 to said individual; and
- (b) changing the temperature of at least a portion of said individual from:
 - (i) a permissive temperature to a restrictive temperature, or
 - (ii) a restrictive temperature to a permissive temperature.

27. The method of claim 26 wherein said individual is a human.

28. A method for regulating the expression of a protein or untranslated RNA molecule in an individual comprising:

- (a) administering a recombinant host cell comprising at least one DNA molecule of claim 1 to said individual; and
- (b) changing the temperature of at least a portion of said individual from:
 - (i) a permissive temperature to a restrictive temperature, or
 - (ii) a restrictive temperature to a permissive temperature.

-63-

29. The method of claim 28, wherein said recombinant host cells are obtained from the same individual into which said host cells are administered.

30. The method of claim 29, wherein said recombinant host cells are keratinocytes, epithelial cells, or fibroblasts.

5 31. The method of claim 28, wherein said individual is a human.

32. A pharmaceutical composition comprising at least one DNA molecule of claim 1 and a pharmaceutically acceptable carrier.

33. A pharmaceutical composition comprising at least one RNA molecule of claim 12 and a pharmaceutically acceptable carrier.

10 34. A pharmaceutical composition comprising at least one alphaviral particle of claim 13 and a pharmaceutically acceptable carrier.

35. A genetically engineered, non-human animal having host cells containing at least one DNA molecule of claim 1.

15 36. The animal of claim 35, wherein the DNA molecule is stably integrated into the host cell genome.

37. A genetically engineered, non-human animal having host cells containing at least one RNA molecule of claim 12.

38. A DNA vector system comprising one or more polynucleotides which encode RNA molecules, said RNA molecules comprising:

- 20 (a) at least one *cis*-acting sequence element,
- (b) a first open reading frame having a nucleotide sequence encoding a non-cytopathic, temperature-sensitive RNA-dependent RNA polymerase, and

-64-

(c) at least one second nucleotide sequence selected from the group consisting of:

(i) a second open reading frame encoding a protein, or portion thereof, wherein said second open reading frame is in a translatable format after one or more RNA-dependent RNA replication events;

(ii) a sequence complementary to all or part of the second open reading frame of (i); and

(iii) a sequence encoding an untranslated RNA molecule, or complement thereof.

39. A DNA vector system of claim 38, wherein the RNA-dependent RNA polymerase is of viral origin.

40. A DNA vector system of claim 38, wherein the RNA-dependent RNA polymerase is of alphaviral origin.

41. A DNA vector system of claim 38 which encodes an RNA-dependent RNA polymerase that has replicase activity at temperatures below 34°C and low or undetectable replicase activity at 34°C and above.

42. A DNA vector system of claim 38, wherein the second open reading frame of 1(c)(i) encodes a cytokine, lymphokine, tumor necrosis factor, interferon, toxic protein, prodrug converting enzyme, or other protein.

43. A DNA vector system of claim 38, wherein the second open reading frame of 1(c)(i) encodes human erythropoietin or human β -interferon.

44. A DNA vector system of claim 38, wherein the second nucleotide sequence of 1(c)(ii) or 1(c)(iii) encodes an untranslated RNA molecule selected from the group consisting of an antisense RNA molecule, tRNA molecule, rRNA molecule, ribozyme.

-65-

45. A method of making a recombinant host cell comprising introducing at least one polynucleotide of claim 38 into a host cell.

46. An *in vitro* cell culture comprising a recombinant host cell produced by the method of claim 45.

5 47. An *in vitro* cell culture comprising a recombinant host cell comprising at least one polynucleotide of claim 38.

48. The cell culture of claim 47, wherein some or all of the polynucleotide sequences of claim 38 are stably maintained in said host cell.

10 49. A composition comprising one or more RNA molecules transcribed from one or more polynucleotides of the vector system of claim 38.

50. An alphaviral particle containing at least one RNA molecule of claim 49.

51. An *in vitro* cell culture comprising a recombinant host cell comprising at least one RNA molecule of claim 49.

15 52. A method for producing a protein or untranslated RNA molecule in a recombinant host cell comprising:

- (a) growing host cells under suitable culture conditions;
- (b) introducing at least one polynucleotide of claim 38 into said host cells; and
- 20 (c) recovering said protein or untranslated RNA molecule.

53. A method for producing a protein or untranslated RNA molecule in a recombinant host cell comprising:

- (a) growing host cells under suitable culture conditions;
- (b) introducing at least one RNA molecule of claim 49 into

-66-

said host cells; and

- (c) recovering said protein or untranslated RNA molecule.

54. The method of claim 53, wherein the protein is erythropoietin.

55. The method of claim 53, wherein said RNA is packaged into an alphaviral particle.

56. A method for producing alphaviral particles containing an RNA molecule of claim 50 comprising:

- (a) growing host cells under suitable culture conditions;
- (b) introducing into said host cells at least one polynucleotide

of claim 38 having one or more open reading frames which encode alphaviral structural proteins; and

- (c) recovering said alphaviral particles.

57. A method for producing a protein in a recombinant host cell comprising:

- (a) growing host cells under suitable culture conditions;
- (b) infecting said host cells with alphaviral particles produced

by the method of claim 56; and

- (c) recovering said protein.

58. The method of claim 57, wherein said protein is erythropoietin.

59. A method for regulating the expression of a protein or untranslated RNA molecule in a recombinant host cell comprising:

- (a) growing host cells under suitable culture conditions;
- (b) introducing at least one polynucleotide of claim 38 into

said host cells; and

- (c) changing the temperature of the host cell culture from:
 - (i) a permissive temperature to a restrictive

-67-

temperature. or

(ii) a restrictive temperature to a permissive temperature.

5 60. A method for regulating the expression of a protein or untranslated RNA molecule in a recombinant host cell comprising:

(a) growing host cells under suitable culture conditions;

(b) introducing at least one RNA molecule of claim 49 into said host cells; and

(c) changing the temperature of the host cell culture from:

10 (i) a permissive temperature to a restrictive temperature, or

(ii) a restrictive temperature to a permissive temperature.

15 61. A method for regulating the expression of a protein or untranslated RNA molecule in an individual comprising:

(a) administering at least one polynucleotide of claim 38 to said individual; and

(b) changing the temperature of at least a portion of said individual from:

20 (i) a permissive temperature to a restrictive temperature, or

(ii) a restrictive temperature to a permissive temperature.

25 62. A method for regulating the expression of a protein or untranslated RNA molecule in an individual comprising:

(a) administering at least one RNA molecule of claim 49 to said individual; and

(b) changing the temperature of at least a portion of said individual from:

-68-

(i) a permissive temperature to a restrictive temperature, or

(ii) a restrictive temperature to a permissive temperature.

5 63. The method of claim 62, wherein said individual is a human.

64. A method for regulating the expression of a protein or untranslated RNA molecule in an individual comprising:

(a) administering a recombinant host cell comprising at least one polynucleotide of claim 38 to said individual; and

10 (b) changing the temperature of at least a portion of said individual from:

(i) a permissive temperature to a restrictive temperature, or

15 (ii) a restrictive temperature to a permissive temperature.

65. The method of claim 64, wherein said recombinant host cells are obtained from the same individual into which said host cells are administered.

66. The method of claim 65, wherein said recombinant host cells are keratinocytes, epithelial cells, or fibroblasts.

20 67. The method of claim 64, wherein said individual is a human.

68. A pharmaceutical composition comprising at least one polynucleotide of claim 38 and a pharmaceutically acceptable carrier.

69. A pharmaceutical composition comprising at least one RNA molecule of claim 49 and a pharmaceutically acceptable carrier.

-69-

70. A pharmaceutical composition comprising at least one alphaviral particle of claim 50 and a pharmaceutically acceptable carrier.

71. A genetically engineered, non-human animal having host cells containing at least one polynucleotide of claim 38.

5 72. The animal of claim 71, wherein the polynucleotide is stably integrated into the host cell genome.

73. A genetically engineered, non-human animal having host cells containing at least one RNA molecule of claim 49.

10 74. A composition comprising one or more RNA molecules, said RNA molecules comprising:

- (a) at least one *cis*-acting sequence element,
- (b) a first open reading frame having a nucleotide sequence encoding a non-cytopathic, temperature-sensitive RNA-dependent RNA polymerase, and
- 15 (c) at least one second nucleotide sequence selected from the group consisting of:
 - (i) a second open reading frame encoding a protein, or portion thereof, wherein said second open reading frame is in a translatable format after one or more RNA-dependent RNA replication events;
 - 20 (ii) a sequence complementary to all or part of the second open reading frame of (i); and
 - (iii) a sequence encoding an untranslated RNA molecule, or complement thereof.

1/17




FIG.1

2/17




FIG.2

3/17

CTGACGCGCCCTGTAGCGGCGCATTAAAGCGCGGCGGGTGTGGTGGTTACGCGCAGCGTGACCGCTACACTTGCCAGCGCC
CTAGCGCCCGCTCCTTTGCTTTCTCCCTTCTTTCTGCCACGTTGCGCGGCTTTCCCGTCAAGCTCTAAATCGGGG
GCTCCCTTTAGGGTTCCGATTTAGTGCTTTACGGCACCTCGACCCAAAAAACTTGATTAGGGTGATGGTTCACGTAGTG
GGCCATCGCCCTGATAGACGGTTTTTCGCCCTTTGACGTTGGAGTCCACGTTCTTTAATAGTGGACTCTTGTTCCAACT
GGAACAACACTCAACCCTATCTCGGTCTATTCTTTGATTATAAGGGATTTTGCCGATTTGCGCCTATTGGTTAAAAAA
TGAGCTGATTTAACAAAAATTTAACGCGAATTTAACAAAATATTACGCTTACAATTTCCATTGCCATTACAGGCTGCG
CAACTGTTGGGAAGGGCGATCGGTGCGGGCCTTTCGCTATTACGCCAGCTGGCGAAAGGGGGATGTGCTGCAAGGCGAT
TAAGTTGGGTAAACCCAGGGTTTTCCAGTCACGACGTTGTAAAACGACGGCCAGTGAgcgcgcaattaaccctcactaa
agggaaacaaaagctggctagtGATCCAGTCTTATGCAATACTCTTGAGTCTTGCAACATGGTAACGATGAGTTAGCAA
CATGCCCTTACAAGGAGAGAAAAAGCACCGTGATGCCGATTGGTGGAAGTAAGGTGGTACGATCGTGCCTTATTAGGAAG
GCAACAGACGGGTCTGACATGGATTGGACGAACCACTGAATTCGCATTGCAGAGATATTGTATTTAAGTGCCTACCTc
gataccgTCGAGATTGACGGCGTAGTACACACTATTGAATCAAACAGCCGACCAATTGCACTACCATCACAATGGAGAAG
CCAGTAGTAAACGTAGACGTAGACCCCGAGTCCGTTTGTGCTGCAACTGCAAAAAAGCTTCCCGCAATTTGAGGTAGT
AGCACAGCAGGTCACTCCAAATGACCATGCTAATGCCAGAGCATTTTCGCATCTGGCCAGTAACTAATCGAGCTGGAGG
TTCTTACCACAGCGACGATCTTGACATAGGCAGCGCACCGGCTCGTAGAATGTTTTCCGAGCACCAGTATCATTGTGTC
TGCCCCATGCGTAGTCCAGAAGACCCGGACCGCATGATGAAATACGCCAGTAACTGGCGGAAAAAGCGTGCAAGATTAC
AAACAAGAACTTGCATGAGAAGATTAAGGATCTCCGGACCGTACTTGATACGCCGGATGCTGAAACACCATCGCTCTGCT
TTCACAACGATGTTACCTGCAACATGCGTGCCGAATATTCCGTCATGCAGGACGTGTATATCAACGCTCCCGGAATATC
TATCATCAGGCTATGAAAGGCGTGCGGACCCTGTACTGGATTGGCTTCGACACCACCCAGTTCATGTTCTCGGCTATGGC
AGGTTGCTACCTGCGTACAACCAACTGGGCCGACGAGAAAGTCTTGAAGCGCGTAACATCGGACTTTGCAGCACAA
AGCTGAGTGAAGGTAGGACAGGAAAATTGTGATAATGAGGAAGAAGGAGTTGAAGCCCGGGTCGCGGGTTTTATTCTCC
GTAGGATCGACACTTTATCCAGAACACAGAGCCAGCTTGCGAGAGCTGGCATCTTCCATCGGTGTTCCACTTGAATGGAAA
GCAGTCGTACACTTGCCGCTGTGATACAGTGGTGAGTTGCGAAGGCTACGTAGTGAAGAAAAATCACCATCAGTCCCGGA
TCACGGGAGAAACCGTGGGATACGCGGTTACACACAATAGCGAGGGCTTCTTGCTATGCAAAGTTACTGACACAGTAAAA
GGAGAACGGGTATCGTTCCTGTGTGCACGTACATCCCGCCACCATATGCGATCAGATGACTGGTATAATGGCCACGGA
TATATCACCTGACGATGCACAAAACTTCTGGTTGGGCTCAACCAGCGAATTGTCATTAAACGGTAGGACTAACAGGAACA
CCAAACCATGCAAAATTACCTTCTGCCGATCATAGCACAAGGGTTCAGCAAATGGGCTAAGGAGCGCAAGGATGATCTT
GATAACGAGAAAAATGCTGGGTACTAGAGAACGCAAGCTTACGTATGGCTGCTTGTTGGGCGTTTCGCACTAAGAAAGTACA
TTCGTTTTATCGCCACCTGGAACGACAGCTGCGTAAAAGTCCCAGCCTCTTTAGCGCTTTTCCATGTGCTCCGTAT
GGACGACCTCTTTGCCATGTGCTGAGGCAGAAATTGAACTGGCATTGCAACCAAAGAAGGAGGAAAAACTGCTGCAG
GTCTCGGAGGAATTAGTCATGGAGGCCAAGGCTGCTTTTGAGGATGCTCAGGAGGAAGCCAGAGCGGAGAAGCTCCGAGA
AGCACTTCCACCATTAGTGGCAGACAAAGGCATCGAGGCAGCCGAGAAGTTGTCTGCGAAGTGGAGGGGCTCCAGGCGG
ACATCGGAGCAGCATTAGTTGAAACCCGCGCGGTACGTAAGGATAATACCTCAAGCAAATGACCGTATGATCGGACAG
TATATCGTTGTCTCGCCAACTCTGTGCTGAAGAATGCCAACTCGCACCAGCGCACCCGCTAGCAGATCAGGTTAAGAT
CATAACACACTCCGAAGATCAGGAAGGTACGCGGTGCAACCATACGACGCTAAAGTACTGATGCCAGCAGGAGGTGCCG
TACCATGGCCAGAATTCCTAGCACTGAGTGAGAGCGCCACGTTAGTGTACAACGAAAGAGAGTTTGTAACCGCAAATA
TACCATTGGCATGTCATGGCCCCGCAAGAATACAGAAGAGGAGCAGTACAAGGTTACAAAGGCAGAGCTTGCGAAAC
AGAGTACGTGTTTGACGTGGACAAGAAGCGTTGCGTTAAGAAGGAAGAAGCCTCAGGTCTGGTCTCTCGGGAGAACTGA
CCAACCTCCCTATCATGAGCTAGCTCTGGAGGGACTGAAGACCCGACCTGCGGTCCCGTACAAGGTGCAACCAATAGGA
GTGATAGGCACACCGGGTTCGGCAAGTCAGCTATTATCAAGTCAACTGTACGGCACGAGATCTTGTTACCAGCGGAAA
GAAAGAAAATTGTGCGGAAATTGAGGCCGACGTGCTAAGACTGAGGGGTATGCAGATTACGTGGAAGACAGTAGATTGCG
TTATGCTCAACGGATGCCACAAGCCGTAGAAGTGCTGACGTTGACGAAGCGTTGCGGTGCCACGCGAGGAGCACTACTT
GCCCTTGATTGCTATCGTCAGGCCCCGCAAGAAGGTAGTACTATGCGGAGACCCATGCAATGCGGATTCTTCAACATGAT
GCAACTAAAGGTACATTTCAATCACCTGAAAAAGACATATGCACCAAGACATTCTACAAGTATATCTCCCGCGTTGCA

FIG.3A

4/17

CACAGCCAGTTACAGCTATTGTATCGACACTGCATTACGATGGAAGATGAAAACCACGAACCCGTGCAAGAAGAACATT
GAAATCGATATTACAGGGGCCACAAAGCCGAAGCCAGGGGATATCATCTGACATGTTCCGCGGGTGGGTTAAGCAATT
GCAAATCGACTATCCCGGACATGAAGTAATGACAGCCGCGCCTCACAAGGGCTAACAGAAAAGGAGTGTATGCCGTCC
GGCAAAAAGTCAATGAAAACCCACTGTACCGGATCACATCAGAGCATGTGAACGTGTTGCTCACCCGCACTGAGGACAGG
CTAGTGTGAAAAACCTTGACAGGGCGACCCATGGATTAAGCAGCCCACTAACATACCTAAAGGAACTTTTCAGGCTACTAT
AGAGGACTGGGAAGCTGAACACAAGGGAATAATTGCTGCAATAAACAGCCCCACTCCCCGTGCCAATCCGTTTCAGCTGCA
AGACCAACGTTTGTGCGGCGAAAGCATTGGAACCGATACTAGCCACGGCCGGTATCGTACTTACCGGTTGCCAGTGGAGC
GAACTGTTCCACAGTTTTCGGATGACAAACCACATTTCGGCCATTTACGCCCTTAGACGTAATTTGCATTAAGTTTTTCGG
CATGGACTTGACAAGCGGACTGTTTTCTAAACAGAGCATCCCACTAACGTACCATCCCGCGATTTCAGCGAGGCCGGTAG
CTCATTGGGACAACAGCCAGGAACCCGCAAGTATGGGTACGATCACGCCATTGCCGCGAACTCTCCCGTAGATTTCCG
GTGTTCCAGCTAGCTGGGAAGGGCACAACTTGATTTGCAGACGGGGAGAACCAGAGTTATCTCTGCACAGCATAACCT
GGTCCCGGTGAACCGCAATCTTCTCAGCCCTTAGTCCCCGAGTACAAGGAGAAGCAACCCGGCCCGGTCAAAAAATTCT
TGAACCAGTTCAAAACCACTCAGTACTTGTGGTATCAGAGGAAAAATTGAAGCTCCCCGTAAGAGAATCGAATGGATC
GCCCCGATTGGCATAGCCGGTGCAGATAAGAACTACAACCTGGCTTTTCGGGTTTCCGCCGAGGCACGGTACGACCTGGT
GTTTCATCAACATTGGAATAAATACAGAAACCACCACTTTTCAGCAGTGCGAAGACCATGCGGCGACCTTAAAAACCTTT
CGCGTTCCGGCCCTgaattgTTtAaAcTcaggaggcacCCTCGTGGTGAAGTCTTATGGCTACGCCGACCGCAACAGTGAG
GACGTAGTCACCGCTCTTGCCAGAAAGTTTGTTCAGGGTGTCTGCAGCGAGACCAGATTGTGTCTCAAGCAATACAGAAAT
GTACCTGATTTTCCGACAACCTAGACAACAGCCGTACACGGCAATTACCCCCGACCATCTGAATTGCGTGATTTTCGTCCG
TGTATGAGGGTACAAGAGATGGAGTTGGAGCCGCGCCGTACATCCGCAACAAAAGGGAGAATATTGCTGACTGTCAAGAG
GAAGCAGTTGTCAACGCAGCAATCCGTGGGTAGACCAGGCGAAGGAGTCTGCCGTGCCATCTATAACGTTGGCCGAC
CAGTTTTACCGATTTCAGCCACGGAGACAGGCACCGCAAGAATGACTGTGTGCCTAGGAAAGAAAGTGATCCACGCGGTCCG
GCCGTGATTTCCGGAAGCACCCAGAACGAGAAGCCTTGAAATTGCTACAAAACGCCTACCATGCAGTGGCAGACTTAGTA
AATGAACATAACATCAAGTCTGTGCCATTCCACTGCTATCTACAGGCATTTACGCAGCCGAAAAGACCGCCTTGAAGT
ATCATTAACTGCTTGACAACCGCGTAGACAGAACTGACGCGGACGTAACCATCTATTGCCTGGATAAGAAGTGAAGG
AAAGAATCGACGCGGCACTCCAACCTAAGGAGTCTGTAAACAGAGCTGAAGGATGAAGATATGGAGATCGACGATGAGTTA
GTATGGATTCATCCAGACAGTTGCTTGAAGGGAAGAAAGGGATTTCAGTACTACAAAAGGAAAATTGTATTTCGTACTTCGA
AGGCACCAAAATTCATCAAGCAGCAAAAGACATGGCGGAGATAAAGGTCTGTTCCCTAATGACCAGGAAAGTAATGAAC
AACTGTGTGCCTACATATTGGGTGAGACCATGGAAGCAATCCGCGAAAAGTGCCCGGTGACCATTAACCCGTGCTTAGC
CCGCCCCAAAACGTTGCCGTGCTTTGCATGTATGCCATGACGCCAGAAAGGTCACAGACTTAGAAGCAATAACGTCAA
AGAAGTTACAGTATGCTCTCCACCCCCCTTCTAAGCACAAAATTAAGAATGTTTCAAGGTTTCAGTGCACGAAAGTAG
TCCTGTTTAATCCGCACACTCCCGCATTGCTTCCCGCCCGTAAGTACATAGAAGTGCCAGAACAGCCTACCGCTCTCTCT
GCACAGGCCGAGGAGGCCCCGAAGTTGTAGCGACACCGTCACCATCTACAGCTGATAACACCTCGCTTGATGTACAGA
CATCTCACTGGATATGGATGACAGTAGCGAAGGCTCACTTTTTTCGAGCTTTAGCGGATCGGACAACCTCTATTACTAGTA
TGGACAGTTGGTCGTGAGGACCTAGTTCACTAGAGATAGTAGCCGAAGGCAGGTGGTGGTGGCTGACGTTTCATGCCGT
CAAGAGCCTGCCCCATTTCACCGCCAAGGCTAAAGAAGATGGCCCGCCTGGCAGCGGCAAGAAAAGAGCCCACTCCACC
GGCAAGCAATAGCTCTGAGTCCCTCCACCTCTCTTTTGGTGGGTATCCATGTCCCTCGGATCAATTTTCGACGGAGAGA
CGGCCCGCCAGGCAGCGGTACAACCCCTGGCAACAGGCCCCACGGATGTGCCTATGTCTTTTCGGATCGTTTTTCGACGGA
GAGATTGATGAGCTGAGCCGACAGTAAGTGAAGTCCGAACCCGTCCTGTTTGGATCATTTGAACCGGGCGAAGTGAAGTCT
AATTATATCGTCCCGATCAGCCGTATCTTTTCCACTACGCAAGCAGAGACGTAGACGCAGGAGCAGGAGGACTGAATACT
GACTAACCGGGTAGTGGGTACATATTTTCGACGGACACAGGCCCTGGGCACTTGCAAAAAGAGTCCGTTCTGCAGAAC
CAGCTTACAGAACCGACCTTGGAGCGCAATGTCTTGGAAAGAATTTCATGCCCGGTGCTCGACACGTGAAAGAGGAACA
ACTCAAACCTCAGGTACAGATGATGCCACCGAAGCCAACAAAAGTAGGTACAGTCTCGTAAAGTAGAAAATCAGAAAG
CCATAACCACTGAGCGACTACTGTCAGGACTACGACTGTATAACTCTGCCACAGATCAGCCAGAATGCTATAAGATCACC
TATCCGAAACCATGTACTCCAGTAGCGTACCGGCGAACTACTCCGATCCACAGTTCGCTGTAGCTGTCTGTAAACCTA

FIG. 3B

5/17

TCTGCATGAGAACTATCCGACAGTAGCATCTTATCAGATTACTGACGAGTACGATGCTTACTTGGATATGGTAGACGaGA
CAGTCGCaTGCCTGGATACTGCAACCTTCTGCCCGCTAAGCTTAGAAGTTACCCGAAAAACATGAGTATAGAGCCCCG
AATATCCGCAGTGCGGTTCCATCAGCGATGCAGAACACGCTACAAAATGTGCTCATTGCCGCAACTAAAAGAAATTGCAA
CGTCACGCAGATGCGTGAACGCCAACACTGGACTCAGCGACATTCAATGTCGAATGCTTTCGAAAAATATGCATGTAATG
ACGAGTATTGGGAGGAGTTCGCTCGGAAGCCAATTAGGATTACCACTGAGTTTGTACCGCATATGTAGCTAGACTGAAA
GGCCCTAAGGCCGCGCACTATTTGCAAAGACGTATAATTTGGTCCCATTGCAAGAAGTGCCATGGATAGATTGTCAT
GGACATGAAAAGAGACGTGAAAGTTACACAGGCACGAAACACACAGAAGAAAGACCGAAAGTACAAGTGATAACAAGCCG
CAGAACCCCTGGCGACTGCTTACTTATGCGGGATTACCGGGAATTAGTGCGTAGGCTTACGGCCGTCTTGCTTCCAAAC
ATTACACGCTTTTTGACATGTGCGCGGAGGATTTTGATGCAATCATAGCAGAACACTTCAAGCAAGGCGACCCGGTACT
GGAGACGGATATCGCATCATTCGACAAAAGCCAAGACGACGCTATGGCGTTAACCGGTCTGATGATCTTGGAGGACCTGG
GTGTGGATCAACCACTACTCGACTTGATCGAGTGCGCCTTTGGAGAAATATCATCCACCCATCTACCTACGGGTACTCGT
TTTAAATTGGGGCGATGATGAAATCCGGAATGTTCTCACACTTTTGTCAACACAGTTTGAATGTGTTATCGCCAG
CAGAGTACTAGAAGAGCGGCTTAAACGTCCAGATGTGCAGCGTTTATTGGCGACGACAACATCATACATGGAGTAGTAT
CTGACAAAGAAATGGCTGAGAGGTGCGCCACCTGGCTCAACATGGAGGTTAAGATCATCGACGCAGTCATCGGTGAGAGA
CCACCTTACTTCTGCGCGGATTTATCTTGCAAGATTGCGTTACTTCCACAGCGTGCCGCGTGGCGGATCCCTGAAAAG
GCTGTTTAAGTTGGGTAAACCGCTCCAGCCGACGACGAGCAAGACAGAAGACGCGCTCTGCTAGATGAAACAA
AGGCGTGTTTAGAGTAGGTATAACAGGCACTTTAGCAGTGGCGTGACGACCCGGTATGAGGTAGACAATATTACACCT
GTCCTACTGGCATTGAGAACTTTTGCCAGAGCAAAAGAGCATTCCAAGCCATCAGAGGGGAAATAAAGCATCTCTACGG
TGGTCTAAATAGTCAGCATAGTACATTTATCTGACTAATACTACAACACCACCCTctagaCGCGTAGATctcacgt
gagcatgcaggccttgggCCCAATGATCCGACCAGCAAACTCGATGTACTTCCGAGGAACGATGTGCATAATGCATCA
GGCTGGTACATTAGATCCCGCTTACCGCGGGCAATATAGCAACACTAAAACTCGATGTACTTCCGAGGAAGCGCAGTG
CATAATGCTGCGCAGTGTTGCCACATAACCACTATATTAACCATTTATCTAGCGGACGCCAAAACTCAATGTATTTCTG
AGGAAGCGTGGTGATAATGCCACGCAGCGTCTGCATAACTTTTATTATTTCTTTTATTAATCAACAAAATTTTGT
AACATTTCAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAGGGAATTCCAACCTGTTTATTGCAGCTTATAATG
GTTACAAATAAAGCAATAGCATCACAAATTTACAAATAAAGCATTTTTTCACTGCATTCTAGTTGTGGTTGTCCAAA
CTCATCAATGTATCTTATCATGTCTGGATCCGTCGAGACGCGTccaattcgccctatagtgagtcgtattacgcgcgctT
GGCGTAATCATGGTCATAGCTGTTTTCTGTGTGAAATTGTTATCCGCTCACAATTCACACAACATACGAGCCGGAAGCA
TAAAGTGTAAGCCTGGGGTGCTAATGAGTGAGTAACTCACATTAATTGCGTTGCGCTCACTGCCCGCTTTCCAGTCG
GGAAACCTGTGCTGCCAGCTGCATTAATGAATCGGCCAACGCGCGGGGAGAGCGGTTTTCGCTATTGGGCGCTCTTCCGC
TTCCTCGCTCACTGACTCGTGCCTCGGTGCTTGGCTGCGGCGAGCGGTATCAGCTCACTCAAAGCGGTAAACGGT
TATCCACAGAATCAGGGGATAACGCAGGAAGAACATGTGAGCAAAAGGCCAGCAAAAGGCCAGGAACCGTAAAAAGGCC
GCGTTGCTGGCGTTTTTCCATAGGCTCCGCCCCCTGACGAGCATCACAAAATCGACGCTCAAGTCAGAGGTGGCGAAA
CCCGACAGGACTATAAAGATACCAGGCGTTTCCCCCTGGAAGCTCCCTCGTGCGCTCTCCTGTTCCGACCCTGCCGCTTA
CCGGATACCTGTCCGCTTTCTCCCTTCGGAAGCGTGCGCTTTTCTCAATGCTCAGCTGTAGGTATCTCAGTTCCGGTG
TAGGTGCTTCGCTCCAAGCTGGGCTGTGTGCAGAACCCCCGTTACGCCGACCGCTGCGCCTTATCCGGTAACATATCG
TCTTGAGTCCAACCCGTAAGACACGACTTATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGCGAGGTATG
TAGGCGGTGCTACAGAGTTCTTGAAGTGGTGGCTAACTACGGCTACACTAGAAGGACAGTATTTGGTATCTGCGCTCTG
CTGAAGCCAGTTACCTTCGGAAGAGTTGGTAGCTCTTGATCCGGCAACAAACCACCGCTGGTAGCGGTGGTTTTTT
TGTTTGCAAGCAGCAGATTACGCGCAGAAAAAAGGATCTCAAGAAGATCCTTTGATCTTTTCTACGGGTCTGACGCTC
AGTGAACGAAAACTCACGTTAAGGGATTTTGGTCATGAGATTACAAAAAGGATCTTACCTAGATCCTTTTAAATTAA
AAATGAAGTTTAAATCAATCTAAAGTATATAGAGTAACTTGGTCTGACAGTTACCAATGCTTAATCAGTGAGGCACC
TATCTCAGCGATCTGTCTATTTGTTTCATCCATAGTTGCTGACTCCCGTCGTGTAGATAACTACGATACGGGAGGGCT
TACCATCTGGCCCCAGTGCTGCAATGATACCGCGAGACCCAGCTCACCAGGCTCCAGATTTATCAGCAATAAACCAGCCA
GCCGGAAGGGCCGAGCGCAGAAGTGGTCTTGAACCTTTATCCGCTCCATCCAGTCTATTAATTGTTGCCGGGAAGCTAG

FIG.3C

6/17

AGTAAGTAGTTCGCCAGTTAATAGTTTGCACAACGTTGTTGCCATTGCTACAGGCATCGTGGTGTCACGCTCGTCGTTTG
GTATGGCTTCATT CAGCTCCGGTTCCCAACGATCAAGGCGAGTTACATGATCCCCATGTTGTGCAAAAAAGCGGTAGC
TCCTTCGGTCCTCCGATCGTTGT CAGAAGTAAGTTGGCCGAGTGTTATCACTCATGGTTATGGCAGCACTGCATAATTC
TCTTACTGTCATGCCATCCGTAAGATGCTTTTCTGTGACTGGTGAGTACTCAACCAAGTCATTCTGAGAATAGTGTATGC
GGCGACCGAGTTGCTCTTGCCCGGCGTCAATACGGGATAATACCGCGCCACATAGCAGAACTTTAAAAGTGCTCATCATT
GGAAAACGTTCTTCGGGGCGAAAACCTCAAGGATCTTACCGCTGTTGAGATCCAGTTCGATGTAACCCACTCGTGCACC
CAACTGATCTTCAGCATCTTTTACTTTACCAGCGTTTCTGGGTGAGCAAAAACAGGAAGGCAAAATGCCGAAAAAAGG
GAATAAGGGCGACACGGAAATGTTGAATACTCATACTCTTCCTTTTCAATATTATTGAAGCATTTATCAGGGTTATTGT
CTCATGAGCGGATACATATTGAATGTATTTAGAAAAATAACAAATAGGGGTTCCGCGCACATTTCCCCGAAAAGTGCC
AC

FIG.3D

7/17




FIG.4A




FIG.4B

8/17




FIG.5A




FIG.5B

9/17




FIG. 6A




FIG. 6B

10/17




FIG.7A

11/17




FIG.7B

12/17



FIG.8A

13/17




FIG.8B

14/17

1 2 3 4



FIG.9

15 / 17




FIG. 10

16/17




FIG.11




FIG.12

-1-

SEQUENCE LISTING

<110> Cytos Biotechnology AG
Renner, Wolfgang A.
Nieba, Lars
Boorsma, Marco

<120> Inducible Alphaviral Gene Expression System

<130> 1700.002PC01

<140>

<141>

<150> US 60/079,562

<151> 1998-03-27

<160> 9

<170> PatentIn Ver. 2.0

<210> 1

<211> 11282

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:cDNA

<400> 1

```
ctgacgcgcc ctgtagcgcc gcattaagcg cggcgggtgt ggtggttacg cgcagcgtga 60
ccgctacact tgcacgcgcc ctacgcgccg ctcccttcgc tttcttcctc tcctttctcg 120
ccaactgtcg cggttttccc cgtcaagctc taaatcgggg gctcccttta gggttccgat 180
ttagtgtttt acggcacctc gaccccaaaa aacttgatta ggtgatggtt tcacgtagt 240
ggccatcgcc ctgatagacg gtttttcgcc ctttgacgtt ggagtccacg ttctttaata 300
gtggactctt gttccaaact ggaacaacac tcaaccctat ctcggtctat tcttttgatt 360
tataagggat tttgccgatt tcggcctatt ggttaaaaaa tgagctgatt taacaaaaat 420
ttaacgcgaa ttttaacaaa atattaacgc ttacaatttc cattcgccat tcaggctgcg 480
caactgttgg gaaggcgat cggtgcgggc ctcttcgcta ttacgccagc tggcgaaagg 540
gggatgtgct gcaaggcgat taagtgggt aacgccaggg tttcccagc cagcagttg 600
taaacgcgac gccagtgagc gcgcaattaa ccctcactaa aggaacaaa agctggctag 660
tggatccagt cttatgcaat actctttagt tcttgcaaca tggtaacgat gagttagcaa 720
catgccttac aaggagagaa aaagcaccgt gcatgccgat tggtaggaat aagggtgtac 780
gatcgtgcct tattaggaag gcaacagacg ggtctgacat ggattggacg aaccactgaa 840
ttccgcattg cagagatatt gtatttaagt gccctacctc gataccgtcg agattgacgg 900
cgtagtacac actattgaat caaacagccg accaattgca ctaccatcac aatggagaag 960
ccagtagtaa acgtagacgt agacccccag agtcggtttg tcgtgcaact gcaaaaaagc 1020
ttcccgaat ttgaggtagt agcacagcag gtcactccaa atgaccatgc taatgccaga 1080
gcattttcgc atctggccag taaactaatc gagctggagg ttcctaccac agcgacgac 1140
ttggacatag gcagcgacc ggctcgtaga atgttttccg agcaccagta tcattgtgtc 1200
tgccccatgc gtagtccaga agaccggac cgcgatgatg aatacgccag taaactggcg 1260
gaaaaagcgt gcaagattac aaacaagaac ttgcatgaga agattaagga tctccggacc 1320
```

-2-

gtacttgata cgccggatgc tgaaacacca tcgctctgct ttcacaacga tgttacctgc 1380
 aacatgcgtg ccgaatatcc cgtcatgcag gacgtgtata tcaacgctcc cggaactatc 1440
 tatcatcagg ctatgaaagg cgtgcggacc ctgtactgga ttggcttcga caccacccag 1500
 ttcattgtct cggtatggc aggttcgtac cctgcgtaca acaccaactg ggccgacgag 1560
 aaagtccttg aagcgcgtaa catcggactt tgcagcacia agctgagtga aggtaggaca 1620
 ggaaaattgt cgataatgag gaagaaggag ttgaagcccg ggtcgcgggt ttatttctcc 1680
 gtaggatcga cactttatcc agaacacaga gccagcttgc agagctggca tcttccatcg 1740
 gtgttccact tgaatggaaa gcagtcgtac acttgccgct gtgatacagt ggtgagttgc 1800
 gaaggctacg tagtgaagaa aatcaccatc agtcccggga tcacgggaga aaccgtggga 1860
 tacgcggtta cacacaatag cgagggtctc ttgctatgca aagtactga cacagtaaaa 1920
 ggagaacggg tatcgttccc tgtgtgcacg tacatcccgg ccaccatag cgatcagatg 1980
 actggtataa tggccacgga tatatcacct gacgatgcac aaaaacttct ggttgggctc 2040
 aaccagcgaa ttgtcattaa cggtaggact aacaggaaca ccaacacccat gcaaaattac 2100
 cttctgccga tcatagcaca aggttcagc aaatgggcta aggagcgcaa ggatgatctt 2160
 gataacgaga aaatgctggg tactagagaa cgcaagctta cgtatggctg cttgtgggag 2220
 tttcgacta agaaagtaca ttcgttttat cgcccacctg gaacgcagac ctgcgtaaaa 2280
 gtcccagcct cttttagcgc ttttcccatg tcgtccgtat ggacgacctc tttgcccag 2340
 tcgctgaggc agaaattgaa actggcattg caaccaaaaga aggaggaaaa actgctgcag 2400
 gtctcgagg aattagtcac ggaggccaag gctgcttttg aggatgctca ggagggaaggc 2460
 agagcggaga agtcccgaga agcacttcca ccattagtgg cagacaaagg catcgaggca 2520
 gccgcagaag ttgtctgcga agtggagggg ctccaggcgg acatcgagagc agcattagtt 2580
 gaaaccccg cgggtcacgt aaggataata cctcaagcaa atgaccgtat gatcggacag 2640
 tatatcgttg tctcgccaaa ctctgtgctg aagaatgcca aactcgcacc agcgcacccg 2700
 ctagcagatc aggttaagat cataacacac tccggaagat cagggaaggta cgcggtcgaa 2760
 ccatacgacg cttaaagtact gatgccagca ggaggtgccg taccatggcc agaattccta 2820
 gcactgagtg agagcgccac gttagtgtac aacgaaagag agtttgtgaa ccgcaacta 2880
 taccacattg ccattgcattg ccccgccaag aatacagaag aggagcagta caaggttaca 2940
 aaggcagagc ttgcagaaac agagtacgtg tttgacgtgg acaagaagcg ttgcgttaag 3000
 aagggaagag cctcaggtct ggtcctctcg ggagaactga ccaaccctcc ctatcatgag 3060
 ctactctgag agggactgaa gaccgcacct gcggtcccg acaaggtcga acaaatagga 3120
 gtgataggca caccggggtc gggcaagtca gctattatca agtcaactgt cacggcacga 3180
 gatcttgta ccagcggaag gaaagaaaat tgcgcgaaa ttgaggccga cgtgctaaga 3240
 ctgaggggta tgcagattac gtcgaagaca gtgattcgg ttatgctcaa cggatgccac 3300
 aaagccgtag aagtgtgtga cgttgacgaa gcgttcgctg gccacgcagg agcactactt 3360
 gccttgattg ctactgtcag gcccgcgaag aaggtagtag tatgcggaga ccccatgcaa 3420
 tgcggattct tcaacatgat gcaactaaag gtacatttca atcaccctga aaaagacata 3480
 tgcaccaaga cattctacaa gtatatctcc cggcgttgca cacagccagt tacagctatt 3540
 gtatcgacac tgcattacga tggaaagatg aaaaccacga acccgtgcaa gaagaacatt 3600
 gaaatcgata ttacaggggc cacaagccg aagccagggg atatcatcct gacatgtttc 3660
 cgcgggtggg ttaagcaatt gcaaatcgac tatcccgagc atgaagtaat gacagcccg 3720
 gcctcacaaag ggctaaccag aaaaggagtg tatgcgtcc ggcaaaaagt caatgaaaac 3780
 ccactgtacg cgatcacatc agagcatgtg aacgtgttgc tcaccgcac tgaggacagg 3840
 ctagtgtgga aaaccttgca gggcgaccca tggattaagc agccactaa catacctaaa 3900
 ggaaactttc aggtactat agaggactgg gaagctgaac acaagggaat aattgctgca 3960
 ataaacagcc ccactccccg tgccaatccg ttcagctgca agaccaacgt ttgctgggag 4020
 aaagcattgg aaccgatact agccacggcc ggtatcgtac ttaccggtg ccagtggagc 4080
 gaactgttcc cacagtttgc gtagtgcgaa ccacattcgg ccatttacgc cttagacgta 4140
 atttgatta agtttttcgg catggacttg acaagcgagc tgttttctaa acagagcatc 4200
 ccactaacgt accatcccgc cgattcagcg aggcggtag ctcatggga caacagccca 4260
 ggaacccgca agtatgggta cgatcacgcc attgccgagc aactctcccg tagatttccg 4320
 gtgttccagc tagctgggaa gggcacacaa cttgatttgc agacggggag aaccagagtt 4380
 atctctgcac agcataacct ggtcccggg aaccgcaatc ttcctcacgc cttagtcccc 4440

gagtacaagg agaagcaacc cggcccggtc aaaaaattct tgaaccagtt caaacaccac 4500
 tcagtacttg tggatcaga ggaaaaaatt gaagctcccc gtaagagaat cgaatggatc 4560
 gccccgattg gcatagccgg tgcagataag aactacaacc tggctttcgg gtttcgccc 4620
 caggcacggg acgacctggt gttcatcaac attggaacta aatacagaaa ccaccacttt 4680
 cagcagtgcg aagaccatgc ggcgacctta aaaacccttt cgcgttcggc cctgaattgt 4740
 ttaaaactcag gaggcacctt cgtggtgaag tcctatggct acgccgaccg caacagttag 4800
 gacgtagtca ccgctcttgc cagaaagtgt gtcagggtgt ctgcagcgag accagattgt 4860
 gtctcaagca atacagaaat gtacctgatt ttccgacaac tagacaacag ccgtacacgg 4920
 caattcacc cgcaccatct gaattgcgtg atttcgtccg tgtatgaggg tacaagagat 4980
 ggagttggag ccgcgccgtc ataccgcacc aaaagggaga atattgctga ctgtcaagag 5040
 gaagcagttg tcaacgcagc caatccgctg ggtagaccag gcgaaggagt ctgccgtgcc 5100
 atctataaac gttggccgac cagttttacc gattcagcca cggagacagg caccgcaaga 5160
 atgactgtgt gcctagggaa gaaagtgtc cagcggtcgc gccctgattt ccggaagcac 5220
 ccagaagcag aagccttgaa attgctacaa aacgcctacc atgcagtggc agacttagta 5280
 aatgaacata acatcaagtc tgtcgccatt ccactgctat ctacaggcat ttacgcagcc 5340
 ggaaaagacc gccttgaaat atcacttaac tgcttgacaa ccgcgctaga cagaactgac 5400
 gcggacgtaa ccatctattg cctggataag aagtggagg aaagaatcga cgcggcactc 5460
 caacttaagg agtctgtaac agagctgaag gatgaagata tggagatcga cgatgagtta 5520
 gtatggattc atccagacag ttgcttgaag ggaagaaagg gattcagtac taaaaagga 5580
 aaattgtatt cgtacttcga aggcaccaa ttccatcaag cagcaaaaga catggcggag 5640
 ataaaggtcc tgttccttaa tgaccaggaa agtaatgaac aactgtgtgc ctacatattg 5700
 ggtgagacca tggaagcaat ccgcgaaaag tgcccggctg accataaccc gtcgtctagc 5760
 ccgcccaaaa cgttgccgtg cctttgcatg tatgccatga cgcagaaaag ggtccacaga 5820
 cttagaagca ataacgtcaa agaagttaca gtatgtcctt ccacccccct tccaaagcac 5880
 aaaattaaga atgttcagaa ggttcagtgc acgaaagtag tccgttttaa tccgcacact 5940
 cccgcattcg ttcccgcggc taagtacata gaagtgccag aacagcctac cgctcctcct 6000
 gcacaggccg aggaggcccc cgaagttgta gcgacaccgt caccatctac agctgataac 6060
 acctcgcttg atgtcacaga catctcactg gatatggatg acagtacgga aggtcactt 6120
 ttttcgagct ttagcggatc ggacaactct attactagta tggacagttg gtcgtcagga 6180
 cctagttcac tagagatagt agaccgaagg caggtggtgg tggtgacgt tcatgccgtc 6240
 caagagcctg cccctattcc accgccaaag ctaaaagaaga tggccgcctt ggcagcgga 6300
 agaaaagagc ccactccacc ggcaagcaat agctctgagt cccctcacct ctcttttgg 6360
 ggggtatcca tgtccctcgg atcaattttc gcggagaga cggcccgcca ggcagcgga 6420
 caacccctgg caacaggccc cagggatgtg cctatgtctt tcggatcgtt ttccgacgga 6480
 gagattgatg agctgagccg cagagtaact gagtccgaac ccgtcctgtt tggatcattt 6540
 gaaccgggag aagtgaactc aattatatcg tcccgatcag ccgtatcttt tccactacgc 6600
 aagcagagac gtagacgag gagcaggagg actgaatact gactaaccgg ggtaggtggg 6660
 tacatatttt cgacggacac aggccctggg cacttgcaaa agaagtccgt tctgcagaac 6720
 cagcttacag aaccgacctt ggagcgcaat gtccgtgaaa gaattcatgc cccggtgctc 6780
 gacacgtcga aagaggaaca actcaaaactc aggtaccaga tgatgccac cgaagccaac 6840
 aaaagtaggt accagtctcg taaagtagaa aatcagaaag ccataaccac tgagcgacta 6900
 ctgtcaggac tacgactgta taactctgcc acagatcagc cagaatgcta taagatcacc 6960
 tatccgaaac cattgtactc cagtagcgta ccggcgaaact actccgatcc acagttcgtc 7020
 gttagctgtc gtaacaacta tctgcatgag aactatccga cagtagcatc ttatcagatt 7080
 actgacgagt acgatgctta cttggatatg gtagacgaga cagtcgcatg cctggatact 7140
 gcaaccttct gcccgctaa gcttagaagt taccgaaaa aacatgagta tagagccccg 7200
 aatatccgca gtgcggttcc atcagcgatg cagaacacgc taaaaatgt gctcattgcc 7260
 gcaactaaaa gaaattgcaa cgtcacgcag atgcgtgaac tgccaacact ggactcagcg 7320
 acattcaatg tcgaatgctt tcgaaaatat gcatgtaatg acgagtattg ggaggagttc 7380
 gctcgggaagc caattaggat taccactgag tttgtaccg catatgtagc tagactgaaa 7440
 ggccctaagg ccgccgact atttgcaaa acgtataatt tggctccatt gcaagaagtg 7500
 cctatggata gattcgtcat ggacatgaaa agagacgtga aagttacacc aggcacgaaa 7560

cacacagaag aaagaccgaa agtacaagtg atacaagccg cagaaccctt ggcgactgct 7620
 tacttatgcg ggattcaccg ggaattagtg cgtaggctta cggccgtctt gcttccaaac 7680
 attcacacgc tttttgacat gtcggcgag gattttgatg caatcatagc agaacacttc 7740
 aagcaaggcg acccggtact ggagacggat atcgcatcat tgcacaaaag ccaagacgac 7800
 gctatggcgt taaccggtct gatgatcttg gaggacctgg gtgtggatca accactactc 7860
 gacttgatcg agtgcgcctt tggagaaata tcatccaccc atctacctac ggttactcgt 7920
 tttaaattcg gggcgatgat gaaatccgga atgttcctca cactttttgt caacacagtt 7980
 ttgaatgctg ttatcgccag cagagtacta gaagagcggc ttaaaacgtc cagatgtgca 8040
 gcgttcattg gcgacgacaa catcatacat ggagtagtat ctgacaaaga aatggctgag 8100
 aggtgcgcca cctgggtcaa catggagggt aagatcatcg acgcagtcac cggtagagaga 8160
 ccaccttact tctgcggcgg atttatcttg caagattcgg ttacttccac agcgtgccgc 8220
 gtggcggatc ccctgaaaag gctgtttaag ttgggtaaac cgctccagc cgacgacgag 8280
 caagacgaag acagaagacg cgctctgcta gatgaaacaa aggcgtggtt tagagtaggt 8340
 ataacaggca ctttagcagt ggccgtgacg acccggtatg aggtagacaa tattacacct 8400
 gtcctactgg cattgagaac ttttgcccag agcaaaagag cattccaagc catcagaggg 8460
 gaaataaagc atctctacgg tggtcctaaa tagtcagcat agtacatttc atctgactaa 8520
 tactacaaca ccaccacctc tagacgcgta gatctcacgt gagcatgcag gccttgggccc 8580
 caatgatccg accagcaaaa ctcgatgtac ttccgaggaa ctgatgtgca taatgcatca 8640
 ggctgtgaca ttagatcccc gcttaccgcg ggcaatatag caaactaaa aactcgatgt 8700
 acttccgagg aagcgcagtg cataatgctg cgcagtggtg ccacataacc actatattaa 8760
 ccatttatct agcggacgcc aaaaactcaa tgtatttctg aggaagcgtg gtgcataatg 8820
 ccacgcagcg tctgcataac ttttattatt tcttttatta atcaacaaa ttttgtttt 8880
 aacatttcaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaggga tcccaactt 8940
 gtttattgca gcttataatg gttacaaata aagcaatagc atcacaaatt tcacaaataa 9000
 agcatttttt tcactgcatt ctagtgtggt tttgtccaaa ctcatcaatg tatcttatca 9060
 tgtctggatc cgtcgagacg cgtccaattc gccctatagt gagtcgtatt acgcgcgctt 9120
 ggcgtaatca tggatcatagc tggttcctgt gtgaaattgt tatccgctca caattccaca 9180
 caacatacga gccggaagca taaagtgtaa agcctggggt gcctaagtag tgagctaact 9240
 cacattaatt gcgttgcgct cactgcccgc tttccagtcg ggaaacctgt cgtgccagct 9300
 gcattaatga atcgcccaac gcgcggggag aggcggtttg cgtattgggc gctcttccgc 9360
 ttctcgcctc actgactcgc tgcgctcgtt cggttcggtg cggcgagcgg tatcagctca 9420
 ctcaaaggcg gtaatacggg tatccacaga atcaggggat aacgcaggaa agaacatgtg 9480
 agcaaaaggc cagcaaaagg ccaggaaccg taaaaggcc gcgttgctgg cgtttttcca 9540
 taggtccgc cccctgacg agcatcacia aaatcgacgc tcaagtcaga ggtggcgaaa 9600
 ccgcagagga ctataaagat accaggcgtt tccccctgga agctccctcg tgcgctctcc 9660
 tgttccgacc ctgcccctta ccggatacct gtccgccttt ctcccttcgg gaagcgtggc 9720
 gctttctcaa tgcacacgt gtaggtatct cagttcgggt taggtcgttc gctccaagct 9780
 gggtgtgtg cacgaacccc ccgttcagcc cgaccgctgc gccttatccg gtaactatcg 9840
 tcttgagtcc aaccgggtaa gacacgactt atcgccactg gcagcagcca ctggtaacag 9900
 gattagcaga gcgaggtatg taggcggtgc tacagagttc ttgaagtgtt ggcctaacta 9960
 cggctacact agaaggacag tatttggtat ctgcgctctg ctgaagccag ttaccttcgg 10020
 aaaaagagtt ggtagctctt gatccggcaa acaaacacc gctggtagcg gtggtttttt 10080
 tgtttgcaag cagcagatta cgcgcagaaa aaaaggatct caagaagatc ctttgcattt 10140
 ttctacgggg tctgacgctc agtgaacga aaactcacgt taagggtatt tggatcatgag 10200
 attatcaaaa aggatcttca cctagatcct tttaaattaa aaatgaagtt ttaaataaat 10260
 ctaaagtata tatgagtaaa ctgtgtctga cagttaccaa tgcttaatca gtgaggcacc 10320
 tatctcagcg atctgtctat ttctgtctat catagttgcc tgactccccg tctgttagat 10380
 aactacgata cgggagggtt taccatctgg cccagtgct gcaatgatac cgcgagaccc 10440
 acgctcaccg gctccagatt tatcagcaat aaaccagcca gccggaaggg ccgagcgag 10500
 aagtggctct gcaactttat ccgcctccat ccagtcctatt aattgttgcc gggaagctag 10560
 agtaagtagt tgcgcagtta atagtttgcg caacgttggt gccattgcta caggcatcgt 10620
 ggtgtcacgc tctgcgtttg gtatggcttc attcagctcc gggtcccaac gatcaaggcg 10680

-5-

```

agttacatga tcccccatgt tgtgcaaaaa agcggttagc tccttcgggc ctccgatcgt 10740
tgtcagaagt aagttggccg cagtgttatc actcatggtt atggcagcac tgcataattc 10800
tcttactgtc atgcatccg taagatgctt ttctgtgact ggtgagtact caaccaagtc 10860
attctgagaa tagtgtatgc ggcgaccgag ttgtcttgc ccggcgtcaa tacgggataa 10920
taccgcgcca catagcagaa ctttaaaagt gctcatcatt ggaaaacgtt cttcggggcg 10980
aaaactctca aggatcttac cgctgttgag atccagttcg atgtaacca ctcgtgcacc 11040
caactgatct tcagcatctt ttactttcac cagcgtttct gggtagcaa aaacaggaag 11100
gcaaaatgcc gcaaaaaagg gaataagggc gacacggaaa tggtgaatac tcatactctt 11160
cctttttcaa tattattgaa gcatttatca gggttattgt ctcagagcg gatacatatt 11220
tgaatgtatt tagaaaaata acaaatagg ggttcgcgc acatttccc gaaaagtgc 11280
ac

```

<210> 2

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 2

aacattgaaa tcgatattac agggg

25

<210> 3

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 3

cgggttatgg tcgaccggc

20

<210> 4

<211> 40

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 4

gtgccctccc ctgagtttaa acaattcagg gccgaacgcg

40

<210> 5

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

-6-

<400> 5
gaattgttta aactcaggag gcaccctcgt gg 32

<210> 6
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:primer

<400> 6
ggtagacgag acagtcgcat gcctggatac 30

<210> 7
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:primer

<400> 7
gtatccaggc atgcgactgt ctcgtctacc 30

<210> 8
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:primer

<400> 8
cagaccgggtt aacgccatag cgtcg 25

<210> 9
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:primer

<400> 9
ctctattact agtatggaca gttgg 25

INTERNATIONAL SEARCH REPORT

International Application No
PCT/IB 99/00523

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/86 C12N9/12 C07K14/18 A61K48/00 A01K67/027
C12N5/10 C12N7/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5 217 879 A (HUANG HENRY V ET AL) 8 June 1993 (1993-06-08) the whole document	1
A	--- STRAUSS, J.H. & STRAUSS, E.G.: "The Alphaviruses: gene expression, replication, and evolution" MICROBIOLOGICAL REVIEWS, vol. 58, no. 3, September 1994 (1994-09), pages 491-562, XP002108834 cited in the application page 498, column 2 - page 499, column 2, paragraph 3; figure 4 page 509, column 2, paragraph 2 - page 517, column 2; figures 14,17 page 525, column 1, paragraph 5 - page 526, column 2, paragraph 3; figure 23 --- -/--	1

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

12 July 1999

Date of mailing of the international search report

27/07/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040. Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Chambonnet, F

INTERNATIONAL SEARCH REPORT

I. International Application No
PCT/IB 99/00523

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	SUOPANKI, J. ET AL.: "Regulation of Alphavirus 26S mRNA transcription by replicase component NSP2" JOURNAL OF GENERAL VIROLOGY., vol. 79, no. 2, February 1998 (1998-02), pages 309-319, XP002108835 SOCIETY FOR GENERAL MICROBIOLOGY, READING., GB ISSN: 0022-1317 the whole document ----	1
A	SHIRAKO, Y. & STRAUSS, J.H.: "Regulation of Sindbis Virus RNA replication : uncleaved P123 and nsP4 function in minus-strand RNA synthesis, whereas cleaved products from P123 are required for efficient plus-strand RNA synthesis" JOURNAL OF VIROLOGY., vol. 68, no. 3, March 1994 (1994-03), pages 1874-1885, XP002108836 ICAN SOCIETY FOR MICROBIOLOGY US the whole document ----	1
A	DE, I. ET AL.: "Sindbis Virus RNA-negative mutants that fail to convert from minus-strand to plus-strand synthesis: role of the nsP2 protein" JOURNAL OF VIROLOGY., vol. 70, no. 5, May 1996 (1996-05), pages 2706-2719, XP002108837 THE AMERICAN SOCIETY FOR MICROBIOLOGY., US ISSN: 0022-538X the whole document ----	1
A	US 5 091 309 A (HUANG HENRY V ET AL) 25 February 1992 (1992-02-25) the whole document ----	1
A	YOUKER, D.R. & SAWICKI S. G.: "Negative strand RNA synthesis by temperature-sensitive mutants of Mouse Hepatitis Virus" ADVANCES IN EXPERIMENTAL MEDECINE AND BIOLOGY, vol. 98, no. 440, 1998, pages 221-226, XP002108838 the whole document ----- -/--	1

INTERNATIONAL SEARCH REPORT

Int. National Application No
PCT/IB 99/00523

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,A	DICOMMO, D. P. & BREMNER R.: "Rapid, high level protein production using DNA-based Semliki Forest virus vectors" JOURNAL OF BIOLOGICAL CHEMISTRY (MICROFILMS), vol. 273, no. 29, 17 July 1998 (1998-07-17), pages 18060-18066, XP002108839 MD US the whole document	1
A	WO 94 17813 A (PARAVAX INC ;GRIEVE ROBERT B (US); XIONG CHENG (US)) 18 August 1994 (1994-08-18) the whole document	1
P,A	WO 98 36779 A (JOHNSTON ROBERT E ;UNIV NORTH CAROLINA (US); DAVIS NANCY L (US); S) 27 August 1998 (1998-08-27) page 11, line 5 - line 29 page 23, line 22 - line 23; claims	1

INTERNATIONAL SEARCH REPORT

international application No.

PCT/IB 99/00523

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 25 to 31, 61 to 67 and partially claims 8 and 45 as far as concerning in vivo gene therapy methods, are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

In International Application No

PCT/IB 99/00523

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5217879 A	08-06-1993	NONE	
US 5091309 A	25-02-1992	NONE	
WO 9417813 A	18-08-1994	AU 6172194 A US 5766602 A	29-08-1994 16-06-1998
WO 9836779 A	27-08-1998	US 5811407 A AU 6168298 A	22-09-1998 09-09-1998